

This article was downloaded by: [de Pietri, Ana Paula Zampieri S]

On: 12 March 2010

Access details: Access Details: [subscription number 919839332]

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Italian Journal of Zoology

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t741771159>

Karyotypes and Ag-NORs in *Phyllomedusa camba* De La Riva, 1999 and *P. rhodei* Mertens, 1926 (Anura, Hylidae, Phyllomedusinae): cytotaxonomic considerations

C. R. Paiva ^a; J. Nascimento ^b; A. P. Z. Silva ^c; P. S. Bernarde ^d; F. Ananias ^a

^a Curso de Ciências Biológicas, Universidade São Francisco (USF), São Paulo, Brazil ^b Curso de Ciências Biológicas, Universidade Braz Cubas (UBC), São Paulo, Brazil ^c Laboratório de Ecologia e Evolução, Instituto Butantan, São Paulo, São Paulo, Brazil ^d Laboratório de Herpetologia, Centro de Ciências Biológicas e da Natureza, Universidade Federal do Acre - UFAC, Campus Floresta, Cruzeiro do Sul, Acre, Brazil

First published on: 10 November 2009

To cite this Article Paiva, C. R., Nascimento, J., Silva, A. P. Z., Bernarde, P. S. and Ananias, F. (2010) 'Karyotypes and Ag-NORs in *Phyllomedusa camba* De La Riva, 1999 and *P. rhodei* Mertens, 1926 (Anura, Hylidae, Phyllomedusinae): cytotaxonomic considerations', Italian Journal of Zoology, 77: 1, 116 – 121, First published on: 10 November 2009 (iFirst)

To link to this Article: DOI: 10.1080/11250000903187585

URL: <http://dx.doi.org/10.1080/11250000903187585>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

SHORT COMMUNICATION

Karyotypes and Ag-NORs in *Phyllomedusa camba* De La Riva, 1999 and *P. rhodei* Mertens, 1926 (Anura, Hylidae, Phyllomedusinae): cytotaxonomic considerations

C. R. PAIVA¹, J. NASCIMENTO², A. P. Z. SILVA³, P. S. BERNARDE⁴, & F. ANANIAS*¹

¹Curso de Ciências Biológicas, Universidade São Francisco (USF), São Paulo, Brazil, ²Curso de Ciências Biológicas, Universidade Braz Cubas (UBC), São Paulo, Brazil, ³Laboratório de Ecologia e Evolução, Instituto Butantan, São Paulo, São Paulo, Brazil, and ⁴Laboratório de Herpetologia, Centro de Ciências Biológicas e da Natureza, Universidade Federal do Acre – UFAC, Campus Floresta, Cruzeiro do Sul, Acre, Brazil

Abstract

The karyotypes of *Phyllomedusa camba* De La Riva, 1999 and *P. rhodei* Mertens, 1926 are presented and the chromosome pairs with Ag-NORs are identified. Both karyotypes have $2n=26$ chromosomes with similar morphology, an exception being the presence of three acrocentric pairs in *P. camba*. In this species the Ag-NORs are found in the proximal region of pairs 1 and 5 whilst in *P. rhodei* an extensive inter-individual variation was observed in the number and position of the Ag-NORs (1p, 3q, 5p, 8p, 11q, and 12q). Based on comparative cytogenetic data of *P. camba* and *P. rhodei*, we discuss the phenetic groups proposed for *Phyllomedusa* genus.

Keywords: Cytogenetic, chromosome, Amphibia, Phyllomedusa, phenetic group

Introduction

The family Hylidae has about 870 species, currently distributed in the subfamilies Hylinae, Pelodyadinae and Phyllomedusinae (Faivovich et al. 2005; Frost 2007). Molecular and morphological studies obtained by Faivovich et al. (2005) show a sister relationship between Pelodyadinae and Phyllomedusinae, which together correspond to the sister taxon of Hylinae. The subfamily Phyllomedusinae is comprised of seven nominal genera: *Agalychnis*, *Hylomantis*, *Cruziophyla*, *Pachymedusa*, *Phasmahyla*, *Phrynomedusa*, and *Phyllomedusa* (Frost 2007). Faivovich et al. (2005) discussed several other characters which are likely to be synapomorphies of Phyllomedusinae, and demonstrated on the basis of molecular data that *Cruziophyla* is the sister taxon of the remaining genera. Two clades were observed: one containing *Phasmahyla* and *Phyllomedusa*, and the other containing *Agalychnis*, *Hylomantis*, *Cruziophyla*, *Pachymedusa* and *Phrynomedusa*. The 30 species of *Phyllomedusa* form a monophyletic group and most of

the species can be distributed amongst five species groups: *burmeisteri*, *hypochondrialis*, *buckeli*, *perinesos* and *tarsius* (Faivovich et al. 2005; Caramaschi 2006; Frost 2007). A few of the Brazilian species were allocated to other genera such as *Hylomanthis*, *Phasmahyla* and *Phrynomedusa* by Cruz (1990). The phylogenetic relationships between the species in these groupings are hitherto not well established.

Phyllomedusa camba De la Riva, 1999 distributed throughout the southeastern Amazon Basin from southwestern Peru, western Brazil (states of Amazon, Acre, and Rondônia) to eastern Bolivia, is included in the *P. tarsius* species group together with *P. boliviana*, *P. coelestis*, *P. tarsius*, *P. trinitatis* and *P. venusta*. Barrio-Amorós (2006) defined the species belonging to the *P. tarsius* group using morphological characters (*P. coelestis*, *P. tarsius*, *P. neildi* sp. nov., *P. trinitatis*, and *P. venusta*) and proposed that *P. boliviana*, *P. camba* and *P. sauvagii* (considered in the group by De la Riva 1999, and Faivovich et al. 2005) should be excluded from the

*Correspondence: F. Ananias, Rua Abílio Ferraro, 237, 13140 000, Paulínia, SP Brazil. Tel: +55 19 3884 7026. Email: feananas@hotmail.com

group because they do not share the most striking feature (herein considered as a synapomorphy) of the group, that is, the golden iris with black reticulations. Although Barrio-Amorós (2006) proposed the exclusion of *P. camba* from the *P. tarsi* species group, its position in the group is maintained (Frost 2007).

Phyllomedusa rohdei Mertens, 1926 is distributed throughout the lowlands of southern Brazil and is included in the *P. hypochondrialis* species group with *P. ayeaye*, *P. azurea*, *P. centralis*, *P. hypochondrialis*, *P. megacephala*, *P. nordestina*, *P. oreades*, and *P. palliata* (Caramaschi 2006; Frost 2007). The species *P. rohdei* has the muscle epicoracoideus, considered a synapomorphy for the group by Faivovich et al. (2005). Although *P. rohdei* was included in the study by Faivovich et al. (2005), the authors did not present a phylogenetic relationship between this species and others in the *P. hypochondrialis* species group, nor with species belonging to other species groups.

With regard to cytogenetic data, most of the information available about the species of *Phyllomedusa* is limited to the diploid number, which is $2n=26$ for most of the species (Beçak et al. 1970; Batistic et al. 1975; Batistic 1989; Kuramoto 1990). Concerning differential staining technique, Batistic (1989) presented C-banding and Ag-NOR data and suggested a hypothesis for the origin of polyploidy in *Phyllomedusa* based on the position of the Ag-NORs. Morando and Hernando (1997) applied the Ag-NOR technique to chromosomes of *P. sauvagii* and *P. hypochondrialis* and detected differences in the number of Ag-NORs between the species as well as a heteromorphism in *P. hypochondrialis*. Kasahara et al. (2007) demonstrated the occurrence of Ag-NORs in *P. distincta* and *P. tetraploidea* and registered the

occurrence of a triploid hybrid. In addition, these authors described the BrdU replication banding pattern in *P. distincta*.

Considering the difficulty in defining the groups of *Phyllomedusa* species as well as the uncertain relationships among species of the *P. tarsi* group, we present for the first time the karyotypes of *P. camba* and *P. rohdei*. Additionally, we also provide Ag-NOR staining for the two species and observe multiple Ag-NORs patterns in the chromosomes of both species.

Materials and methods

Cytogenetic analysis was carried out on two specimens of *Phyllomedusa camba* collected in Ministro Andreazza, state of R ndonia (RO) (11 04'27" S; 61 31'01" W) north of Brazil and on seven specimens of *Phyllomedusa rohdei* collected in Biritiba-Mirim, state of S o Paulo (SP) (23 34'21" S; 46 02'19" W), southeastern Brazil (Figures 1A,B). The voucher specimens were deposited in the Amphibian collection (CFBH) of the Departamento de Zoologia, Instituto de Bioci ncias, UNESP, Rio Claro, SP, Brazil.

The mitotic chromosomes were obtained from direct preparations of bone marrow, liver and testis treated with 0.01% colchicine at a proportion of 0.1 ml/10 g body weight, as described in Baldissera et al. (1993) and Silva et al. (2000), or from the intestine using the technique in Schmid (1978). To improve the mitotic index, we injected phytohemagglutinin in some specimens before the colchicine treatment, at the proportion of 0.1 ml/10 g body weight, 48–72 h before sacrifice. Conventional staining was made with Giemsa 10% diluted in phosphate buffer pH 6.8, and silver nitrate labeling



Figure 1. **A**, adult male of *Phyllomedusa camba* from Ministro Andreazza, R ndonia (RO); **B**, *Phyllomedusa rohdei* from Biritiba-Mirim, S o Paulo (SP).

of the nucleolar organizer regions (Ag-NOR) was obtained by the technique of Howell and Black (1980).

Results and discussion

The species *P. camba* and *P. rhodei* have a diploid number corresponding to $2n=26$, and a karyotype similar to that found for the majority of the Phyllomedusinae species: *Phyllomedusa* (Beçak et al. 1970; Batistic et al. 1975; Batistic 1989; Morando & Hernando 1997; Kasahara et al. 2007); *Agalychnis* (Schmid et al. 1995); and *Pachymedusa* (Schmid 1980). An exception was observed in several tetraploid populations of *P. burmeisteri* from Brazil, with $4n=52$ (Beçak et al. 1970; Batistic et al. 1975; Batistic 1989). The karyotype of *P. rhodei* is made up of metacentric pairs (1, 4, 10, and 13), and submetacentric pairs (2, 3, 5–9, 11 and 12), while in *P. camba*, pairs 1, 4, 7, 10, and 13 are metacentric, 2, 3, 5, 6, and 9 are submetacentric, and 8, 11 and 12 are acrocentric (Figures 2A,B). The occurrence of acrocentric pairs in *P. tarsiuis* observed by Batistic (1989) suggests the proximity of this species and *P. camba*. Nevertheless, these data refute the proposal of Barrio-Amorós (2006), which excluded *P. camba* from the *P. tarsiuis* group. If the presence of acrocentric pairs is considered a synapomorphy, *P. sawagii*, which possesses no acrocentric chromosomes (Batistic 1989), might be excluded from the *P. tarsiuis* group. The karyotype with metacentric and submetacentric chromosomes in *P. rohdei* is compatible to that found by Batistic (1989) for this species and for *P. ayeaye* and *P. hypochondrialis* (all allocated to the *P. hypochondrialis* species group).

As stated by Bogart (1991), centric fusions and fissions are the most likely mechanism for changes in chromosomal number. The subfamily Phyllomedusinae is considered basal in the Hylidae family due to morphological characters and the diploid number of 26 chromosomes (Bogart 1973; Morescalchi 1990). Nevertheless, in the species of *Phyllomedusa* considered to be derived, no alterations have occurred in the number of chromosomes, although the presence of acrocentric chromosomes and polyploidization in some species suggest the presence of speciation events in this anuran group. The species *P. tarsiuis* and *P. camba*, both of them with acrocentric chromosomes, can be considered very closely related and derived in the *Phyllomedusa* genera.

In *P. camba*, 20 metaphases showed Ag-NORs in the proximal regions of the short arms of pair 1, and in the proximal regions of the long arms of pair 5 coincident with secondary constriction (Figure 3B). In *P. rohdei*, the silver staining showed an extensive inter-individual variation in the number and position of Ag-positive regions, in 1p, 3q, 5p, 8p, 11q and 12q (Figure 3A). Inter-individual variation in single or multiple Ag-NOR patterns was described by Wiley et al. (1989) in *Hyla chrysoscelis* and *H. versicolor*, Foote et al. (1991) in *Bufo terrestris*, Miura (1994) in *Rana japonica*, Schmid et al. (1995) in *Agalychnis callidryas*, Kaiser et al. (1996) in *Dendropsophus ebraccatus*, Silva et al. (1999) in *Physalaemus cuvieri*, and Silva et al. (2006) in *Leptodactylus mystacinus*.

In Anura, NOR analysis by silver staining has shown that species, in both primitive and derived families, possess only one pair of Ag-NORs in their diploid karyotype (Schmid 1982; Mahony & Robinson 1986). This observation led King et al. (1990) to suggest the presence of only a single pair of Ag-NORs in diploid karyotypes as an ancestral



Figure 2. **A**, Giemsa-stained karyotypes of *Phyllomedusa rohdei*; **B**, *Phyllomedusa camba*. Note three acrocentric pairs (8, 11 and 12) in *P. camba*.




Figure 3. **A**, Metaphases with Ag-NOR bearing chromosomes of *Phyllomedusa rohdei*; **B**, *Phyllomedusa camba*.

condition in Anura, a hypothesis previously proposed by Schmid (1978) for bufonids and hylids. In the genus *Phyllomedusa*, the occurrence of more than one chromosome with Ag-NORs was observed in *P. burmeisteri*, *Phyllomedusa* sp., *P. distincta*, *P. iheringi*, *P. hypochondrialis*, *P. ayeaye*, *P. tarsiuis*, *P. rohdei* and *P. tetraploidea* (Table I).

Cytochemical tests have revealed that silver does not attach to the ribosomal DNA itself, but rather to proteins specifically associated with this region and to the heterochromatin (Nardi et al. 1978; Sánchez et al. 1995; Dobigny et al. 2002). Nevertheless, in some anuran species, multiple variable Ag-NOR positive region was confirmed using more suitable techniques, such as *in situ* hybridization with a fluorescent rDNA probe (Foote et al. 1991; Kaiser et al. 1996; Lourenço et al. 1998, 2000). These authors suggest that some mechanisms involved in

NOR dispersion in anuran genomes may include inversions and translocations involving chromosomal segments containing NORs, transpositions by mobile genetic elements, amplifications of 'orphan' rDNA cistron, and reinsertion errors during extra-chromosomal amplifications of ribosomal cistrons. Incongruous FISH with rDNA probe data with Ag-staining findings were reported in *Leptodactylus mystacinus* by Silva et al. (2006), and these authors suggested that some Ag-positive sites in the genome of *L. mystacinus* are not true Ag-NORs, but might be related to some peculiarities in the heterochromatin located at these sites, although they are not always identified with C-banding.

The present cytogenetic data are not conclusive for deciding the taxonomic status of *Phyllomedusa* at phenetic species group. Nevertheless, we think that our cytogenetic data are toward a closer proximity

Table I. Multiples Ag-NORs sites in *Phyllomedusa* species.

| Species | Location of the Ag-NORs | Reference |
|-------------------------------------|-------------------------|---------------------------------------|
| <i>Phyllomedusa burmeisteri</i> | 1p 1p 9q 9q | Batistic 1989 |
| <i>Phyllomedusa</i> sp. 4n | 1p 1p 8p 8p | Batistic 1989; Kasahara et al. 2007 |
| <i>Phyllomedusa distincta</i> | 1p 1p 9q 9q | Batistic 1989; Kasahara et al. 2007 |
| <i>Phyllomedusa iheringii</i> | 1p 1p 9q 9q | Batistic 1989 |
| <i>Phyllomedusa sauvagii</i> | 9q 9q | Batistic 1989; Morand & Hernando 1997 |
| <i>Phyllomedusa hypochondrialis</i> | 9–10q 11q | Batistic 1989 |
| | 7p 8q 11q 13q | Morand & Hernando 1997 |
| <i>Phyllomedusa ayeaye</i> | 1p 1p 9q 9q | Batistic 1989 |
| <i>Phyllomedusa tarsiuis</i> | 1p 1p 9q 9q | Batistic 1989 |
| <i>Phyllomedusa camba</i> | 1p 1p 5q 5q | This report |
| <i>Phyllomedusa rohdei</i> | 1p 3q 5p 8p 11q 12q | This report |
| | 2p 8p 8p | Batistic 1989 |

between *P. camba* and *P. tarsi*. Cytogenetic studies of other species of the *P. tarsi* group (sensu De la Riva 1999) are necessary to confirm our hypothesis.

Acknowledgments

We are grateful to Álvaro Dhimas S. Modesto and Marcos Yamamoto for collecting the specimens, Votorantim Celulose e Papel S/A for providing access to animals in the Fazenda São Simão and Instituto Brasileiro do Meio Ambiente e Recursos Naturais Renováveis (IBAMA) for collecting license (02010.005970/03-69). A.P.Z. Silva acknowledges FAPESP for financial support (proc. 2006/06634-8).

References

- Baldissera FA, Oliveira PLS, Kasahara S. 1993. Cytogenetics of four Brazilian *Hyla* species (Amphibia – Anura) and description of a case with a supernumerary chromosome. *Revista Brasileira de Genética* 16:335–345.
- Barrio-Amorós CL. 2006. A new species of *Phyllomedusa* (Anura: Hylidae: Phyllomedusinae) from northwestern Venezuela. *Zootaxa* 1309:55–68.
- Batistic RF. 1989. Aspectos citogenéticos da evolução em *Phyllomedusa* (Anura – Amphibia) [Doctoral Thesis]. Ribeirão Preto (Brazil): Insytuto Butantan. pp 203.
- Batistic RF, Soma M, Beçak ML, Beçak W. 1975. Further studies on polyploidy amphibians. A diploid population of *Phyllomedusa burmeisteri*. *Journal of Heredity* 66:160–162.
- Beçak ML, Denaro L, Beçak W. 1970. Polyploidy and mechanisms of karyotypic diversification in Amphibia. *Cytogenetics* 9:225–238.
- Bogart JP. 1973. Evolution of anuran karyotypes. In: Vial JL, editor. *Evolutionary biology of anurans*. Columbia: University Missouri Press. pp 337–349.
- Bogart JP. 1991. The influence of life history on karyotypic evolution in frogs. In: Green DM, Sessions SK, editors. *Amphibian cytogenetics and evolution*. San Diego, CA: Academic Press. pp 233–257.
- Caramaschi U. 2006. Redefinição do grupo de *Phyllomedusa hypochondrialis*, com redescricao de *P. megacephala* (Miranda-Ribeiro, 1926), revalidação de *P. azurea* Cope, 1862 e descrição de uma nova espécie (Amphibia, Anura, Hylidae). *Arquivos do Museu Nacional do Rio de Janeiro* 64:159–179.
- Cruz CAG. 1990. Sobre as relações intergenéricas de Phyllomedusinae da Floresta Atlântica (Amphibia, Anura, Hylidae). *Revista Brasileira de Biologia* 50:709–726.
- De la Riva I. 1999. A new *Phyllomedusa* from southwestern Amazonia (Amphibia, Anura: Hylidae). *Revista Española de Herpetologia* 13:123–131.
- Dobigny G, Ozouf-Costaz C, Bonillo C, Voloubouev V. 2002. ‘Ag-NORs’ are not always true NORs: New evidence in mammals. *Cytogenetic and Genome Research* 98:75–77.
- Fairovich J, Haddad CFB, Garcia PCA, Frost D, Campbell JA, Wheeler WC. 2005. Systematic review of the frog family Hylidae, with special reference to Hylinae: Phylogenetic analysis and taxonomic revision. *Bulletin of the American Museum of Natural History* 294:1–240.
- Foot DL, Wiley JE, Little ML, Meyne J. 1991. Ribosomal RNA genes site polymorphism in *Bufo terrestris*. *Cytogenetic Cell Genetic* 57:196–199.
- Frost DR. 2007. Amphibian species of the world: An online reference [Internet]. American Museum of Natural History, New York, USA. Available: <http://research.amnh.org/herpetology/amphibia/index.html> (accessed 23 July 2007).
- Howell WM, Black DA. 1980. Controlled silver-staining of nucleolus organizer regions with a protective colloidal developer: A 1-step method. *Experientia* 36:1014–1015.
- Kaiser H, Mais C, Bolanos F, Steinlein C, Feichtinger W, Schmid M. 1996. Chromosomal investigation of three Costa Rican frogs from the 30-chromosome radiation of *Hyla* with the description of a unique geographic variation in nucleolus organizer regions. *Genetica* 98:95–102.
- Kasahara S, Campos JRC, Catroli GF, Haddad CFB. 2007. Cytogenetics of *Phyllomedusa distincta* ($2n=2x=26$), *Phyllomedusa tetraploidea* ($2n=4x=52$) and their triploid hybrids ($2n=3x=39$). 16th International Chromosome Conference, 2007, Amsterdam.
- King M, Contreras N, Honeycutt RL. 1990. Variation within and between nucleolar regions in Australian hylid frogs (Anura) shown by 18s and 28s ‘in-situ’ hybridization. *Genetica* 80:17–29.
- Kuramoto M. 1990. A list of chromosome numbers of anuran amphibians. *Bulletin of Fukuoka University Education* 39:87–127.
- Lourenço LB, Garcia PCA, Recco-Pimentel SM. 2000. Cytogenetics of two species of *Paratelmatobius* (Anura: Leptodactylidae), with phylogenetic comments. *Hereditas* 133:201–209.
- Lourenço LB, Recco-Pimentel SM, Cardoso AJ. 1998. Polymorphism of the nucleolus organizer regions in *Physalaemus petersi* (Amphibia, Anura, Leptodactylidae) detected by silver staining and fluorescence *in situ* hybridization. *Chromosome Research* 6:621–628.
- Mahony MJ, Robinson ES. 1986. Nucleolar organizer region (NOR) location in karyotypes of Australian ground frogs (Family Myobatrachidae). *Genetica* 68:119–127.
- Miura I. 1994. Sex chromosome differentiation in the Japanese Brown frog, *Rana japonica*. II. Sex-linkage analyses of the nucleolar organizer regions in chromosomes no. 4 of the Hiroshima and Saeki populations. *Zoological Science* 11:807–815.
- Morando M, Hernando A. 1997. Localización cromosómica de genes ribosomales activos en *Phyllomedusa hypochondrialis* y *P. sauvagii* (Anura: Hylidae). *Cuadernos Herpetologia* 11:31–36.
- Morescalchi A. 1990. Cytogenetics and the problem of lissamphibian relationships. In: Olmo E, editor. *Cytogenetics of amphibians and reptiles*. Birkhäuser: Verlag. pp 1–19.
- Nardi I, de Lucchini S, Barsacchi-Pilone G, Andronico F. 1978. Chromosome location of the ribosomal RNA genes in *Triturus vulgaris meridionalis* (Amphibia, Urodela). *Chromosoma* 70:91–99.
- Sánchez A, Jimenez R, Burgos M, Stitou S, Zurita F, de la Guardia RD. 1995. Cytogenetic peculiarities in the Algerian hedgehog: Silver stains not only NORs but also heterochromatic blocks. *Heredity* 75:10–16.
- Schmid M. 1978. Chromosome banding in Amphibia. I. Constitutive heterochromatin and nucleolus organizers regions in *Bufo* and *Hyla*. *Chromosoma* 66:361–388.
- Schmid M. 1980. Chromosome banding in Amphibia. IV. Differentiation of GC- and AT-rich chromosome regions in Anura. *Chromosoma* 77:83–103.
- Schmid M. 1982. Chromosome banding in Amphibia VII. Analysis of the structure and variability of NORs in Anura. *Chromosoma* 87:327–344.
- Schmid M, Feichtinger W, Wimer R, Mais C, Bolaños F, Leon P. 1995. Chromosome banding in Amphibia XXI. Inversion

- polymorphism and nucleolus organizer regions in *Agalychnis callidryas* (Anura, Leptodactylidae). *Chromosoma* 101:284–292.
- Silva APZ, Haddad CFB, Galassi GG, Kasahara S. 2006. Multiple nucleolus organizer regions in *Leptodactylus mystacinus* (Amphibia, Anura) and comments on its systematic position in the *L. fuscus* group based on cytogenetic and molecular analyses. *Genetica* 127:35–44.
- Silva APZ, Haddad CFB, Kasahara S. 1999. Nucleolus organizer regions in *Physalaemus cuvieri* (Anura, Leptodactylidae), with evidence of a unique case of Ag-NOR variability. *Hereditas* 131:135–141.
- Silva APZ, Haddad CFB, Kasahara S. 2000. Chromosomal studies on five species of the genus *Leptodactylus* Fitzinger, 1826 (Amphibia, Anura) using differential staining. *Cytobios* 103:25–38.
- Wiley JE, Little ML, Romano MA, Blount DA, Cline GR. 1989. Polymorphism in the location of the 18S and 28S rDNA genes on the chromosomes of the diploid–tetraploid treefrogs *Hyla chrysoscelis* and *Hyla versicolor*. *Chromosoma* 97:481–487.