

NOTES ON GEOGRAPHIC DISTRIBUTION

**Amphibia, Anura, *Barycholos ternetzi*,
Chaunus rubescens, and *Scinax canastrensis*:
Distribution extension, new state record.**

Cybele de Oliveira Araujo¹
Thais Helena Condez²
Célio F. B. Haddad³

¹Seção de Ecologia Florestal, Instituto Florestal.
Rua do Horto 931. CEP 02377-000, São Paulo,
SP, Brazil. E-mail: cyaraujo@iflorestal.sp.gov.br

²Laboratório de Herpetologia, Instituto Butantan.
Avenida Dr. Vital Brazil 1500. CEP 05503-900,
São Paulo, SP, Brazil.

³Departamento de Zoologia, Instituto de
Biociências, Universidade Estadual Paulista.
Caixa Postal 199. CEP 13506-970, Rio Claro, SP,
SP, Brazil.

During field trips on August, September, October 2006, and January 2007, to the Parque Estadual das Furnas do Bom Jesus (PEFBJ) (20°11'14" to 20°16'34" S, 47°22'13" to 47°29'17" W; 647-1018 m), municipality of Pedregulho, northeastern region of the state of São Paulo, southeastern Brazil, we registered three anuran amphibian species representing an increase of their distributions and new records for São Paulo. This state park, with 2069.06 ha, is a representative protected area in the limit of São Paulo and Minas Gerais states. The vegetation of PEFBJ is characterized mostly by secondary seasonal semideciduous forests (Oliveira-Filho and Fontes 2000) along the rivers and on scarps of the canyons and small patches of *Cerrado* (Brazilian savanna; Ratter et al. 1997) in the interior of the park and on the plateau (Branco et al. 1991; Franco et al. 2007).

Barycholos ternetzi (Figure 1) is widely distributed in the Brazilian *Cerrado* domain, being previously known from many localities in the states of Goiás, Mato Grosso, Minas Gerais, Tocantins, Maranhão, and Distrito Federal (Caramaschi and Pombal Jr. 2001; Frost 2007). However, this species has never been recorded in the state of São Paulo. First described as *Paludicola ternetzi* Miranda-Ribeiro, 1937, it was considered a senior synonym of *Barycholos savagei* Lynch, 1980, under the combination *B. ternetzi* (Miranda-Ribeiro, 1937) (Caramaschi and

Pombal Jr. 2001). This small brachycephalid frog (SVL = 25.61 mm, mean weight = 1.75 g, N = 11) is abundant at PEFBJ, where it occurs associated with riverine forest litter and permanent streams with rocky bed in seasonal semideciduous forest and seasonal semideciduous alluvial forest (Franco et al. 2007). We observed many individuals on August and September, but only on October (beginning of the rainy season) this species was found in calling activity.



Figure 1. Adult male of *Barycholos ternetzi* at Parque Estadual das Furnas do Bom Jesus, municipality of Pedregulho, São Paulo, Brazil.

Chaunus rubescens (Figure 2) is known from the states of Pará and Piauí until Goiás and Minas Gerais (Frost 2007). It is a typical species of *Cerrado* areas in central Brazil. This species was first named as *Bufo rufus* Garman, 1877, but this name was considered as a primary homonym of *Bufo rufus* Schneider, 1799 (Lavilla 1992). By this reason the name applied by Lutz (1925) is now in use. Frost et al. (2006) revalidated the generic name *Chaunus* and included *Bufo rubescens* in it, as well as several other species previously included in the genus *Bufo*. In PEFBJ (the first record of this species for São Paulo) this large bufonid toad (SVL males = 84.23 mm, mean weight = 74.54 g, N = 7) was found associated with permanent streams with rocky bed in seasonal semideciduous alluvial forest (Franco et al. 2007) and anthropically disturbed sites, such as artificial permanent ponds. Some individuals were observed on August, September, and October, but only on August (end of dry season) this species was found in calling activity.

NOTES ON GEOGRAPHIC DISTRIBUTION



Figure 2. Adult male of *Chaunus rubescens* at Parque Estadual das Furnas do Bom Jesus, municipality of Pedregulho, São Paulo, Brazil.

Scinax canastrensis (Figure 3), described by Cardoso and Haddad (1982), is known from its type-locality, at Parque Nacional da Serra da Canastra, municipality of São Roque de Minas, state of Minas Gerais (Haddad et al. 1988), and from the municipality of Perdizes, Minas Gerais (Oliveira-Filho and Kokubum 2003). This new record for the state of São Paulo extends the distribution of *S. canastrensis* approximately 45 km to the west of the Parque Nacional da Serra da Canastra, Minas Gerais. This small-size hylid treefrog (SVL males = 29.66 mm, mean weight = 1.92 g, N = 6) occurs at low densities at PEFBJ, where males were observed calling at the end of dry season (August and September) and at the beginning of the rainy season (October) on branches of shrub vegetation, always associated with gallery forest lining the permanent streams in seasonal semideciduous alluvial forest (Franco et al. 2007).

Literature cited

Branco, I. H. D. C., E. N. Domingues, F. C. Sérgio, I. H. Del Cali, I. F. A. Mattos, J. E. A. Berton, M. Rossi, M. R. Eston, R. M. Pfeifer, and W. J. Andrade. 1991. Plano conceitual de manejo - Parque Estadual das Furnas do Bom Jesus, município de Pedregulho, SP. *Revista do Instituto Florestal* 3(2): 137-155.



Figure 3. Adult male of *Scinax canastrensis* at Parque Estadual das Furnas do Bom Jesus, municipality of Pedregulho, São Paulo, Brazil.

The specimens collected were deposited at the Célio F. B. Haddad anuran collection, Departamento de Zoologia, Universidade Estadual Paulista, Rio Claro, São Paulo, Brazil, under the following numbers: *Barycholos ternetzi* (CFBH 13443-51 and 13973), *Chaunus rubescens* (CFBH 13412-13), and *Scinax canastrensis* (CFBH 13438-39 and 13974-78).

Acknowledgements

We thank the Instituto Florestal for financial support; the management officer and staff of Parque Estadual das Furnas do Bom Jesus for field assistance; Cláudia N. Shida for working facilities and field lodging; Flaviana M. de Souza, Ricardo J. Sawaya, and Roque Cielo Filho for very useful comments on the manuscript; and the IBAMA (Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis) for the collection permit. C.F.B.H. thanks FAPESP and CNPq for financial support.

Caramaschi, U. and J. P. Pombal Jr. 2001. *Barycholos savagei*: A junior synonym of *Paludicola ternetzi*, with notes on development. *Journal of Herpetology* 35(2): 357-360.

Cardoso, A. J. and C. F. B. Haddad. 1982. Nova espécie de *Hyla* da Serra da Canastra (Amphibia, Anura, Hylidae). *Revista Brasileira de Biologia* 42(3): 499-503.

NOTES ON GEOGRAPHIC DISTRIBUTION

- Franco, D. C. et al. 2007. Plano de Manejo do Parque Estadual das Furnas do Bom Jesus. Relatório Técnico: Módulo Avaliação Ecológica Rápida - Tema Vegetação. 155 p. Não publicado.
- Frost, D. R. 2007. Amphibian Species of the World: an Online Reference. Version 5.0. Electronic database accessible at <http://research.amnh.org/herpetology/amphibia/index.html>. American Museum of Natural History, New York, USA. Captured on 01 march 2007.
- Frost, D. R., T. Grant, J. Faivovich, R. H. Bain, A. Haas, C. F. B. Haddad, R. O. de Sá, A. Channing, M. Wilkinson, S. C. Donnellan, C. J. Raxworthy, J. A. Campbell, B. L. Blotto, P. Moler, R. C. Drewes, R. A. Nussbaum, J. D. Lynch, D. M. Green, and W. C. Wheeler. 2006. The amphibian tree of life. *Bulletin of the American Museum of Natural History* 297: 1-370.
- Haddad, C. F. B., G. V. Andrade, and A. J. Cardoso. 1988. Anfíbios anuros no Parque Nacional da Serra da Canastra, estado de Minas Gerais. *Brasil Florestal* 64 (24): 9-20.
- Lavilla, E. O. 1992. Tipos portadores de nombres y localidades tipo de anfibios de Argentina. *Acta Zoologica Lilloana* 42: 61-100.
- Lutz, A. 1925. Batraciens du Brésil. *Comptes Rendu des Séances de la Société de Biologie et ses Filiales* 93: 211-214.
- Oliveira-Filho, A. T. and M. A. L. Fontes. 2000. Patterns of floristic differentiation among atlantic forests in Southeastern Brazil and the influence of climate. *Biotropica* 32(4b): 793-810.
- Oliveira-Filho, J. C. and M. N. C. Kokubum. 2003. *Scinax canastrensis*. Brazil: Minas Gerais. *Herpetological Review* 34: 163.
- Ratter, J. A., J. F. Ribeiro, and S. Bridgewater. 1997. The Brazilian cerrado vegetation and threats to its biodiversity. *Annals of Botany* 80: 223-230.

Received April 2007

Accepted May 2007

Published online June 2007