



Phylogenetic relationships and karyotype evolution in the sigmodontine rodent *Akodon* ($2n = 10$ and $2n = 16$) from Brazil

Maria José de J. Silva^{1,2}, James L. Patton³ and Yatiyo Yonenaga-Yassuda¹

¹*Departamento de Biologia, Instituto de Biociências, Universidade de São Paulo, São Paulo, SP, Brazil.*

²*Laboratório de Genética, Instituto Butantan, São Paulo, SP, Brazil.*

³*Museum of Vertebrate Zoology, University of California, Berkeley, CA, USA.*

Abstract

Comparison of G-bands from $2n = 10$ and $2n = 16$ karyotypes of *Akodon* revealed tandem fusions, pericentric inversions and Robertsonian rearrangement in autosomes and addition/deletion of constitutive heterochromatin in sex chromosomes. Cytochrome-b sequences indicate that the $2n = 10$ karyotype is a new species and show it to be a sister taxon of the $2n = 14$, $2n = 15$ and $2n = 16$ karyotypes. Indeed, this group shows a particular evolutionary situation in which a unique taxonomic unit based on morphological data can be detected, but, karyologically, it can be separated into two groups ($2n = 14$ - 15 - 16 and $2n = 10$). Cytochrome-b sequences show a finer resolution, indicating that these four karyotypes represent three molecular entities ($2n = 14$ - 15 , $2n = 16$ and $2n = 10$) that may be derived from a common ancestor with a $2n = 16$ karyotype.

Key words: *Akodon*, cytochrome-b, G-bands, karyotypical evolution, phylogeny.

Received: July 28, 2005; Accepted: December 29, 2005.

Introduction

The Akodontini tribe encompasses about 35% of the total diversity of the extant species among 10 tribes of the subfamily Sigmodontinae of South American rodents (Smith and Patton 1993), although relationships at genera and species levels within this tribe remain uncertain.

Karyotypes have been useful in generating species classification and clarifying some systematic problems, especially those related to cryptic species within the genus *Akodon*. In *Akodon* the diploid chromosome number ranges from $2n = 9$ or 10 to $2n = 52$, with the majority of the members of this genus being characterized by relatively low chromosome numbers (Spotorno and Fernandez, 1976; Yonenaga *et al.*, 1976; Kasahara and Yonenaga-Yassuda, 1984; Liascovich and Reig, 1989; Svartman and Almeida, 1994; Geise *et al.*, 1998; Fagundes and Yonenaga-Yassuda, 1998; Fagundes *et al.*, 1998; Silva and Yonenaga-Yassuda, 1998). Brazilian *Akodon* species with known karyotypes are as follows: *Akodon* sp. n. ($2n = 9, 10$); *A. cursor* (sensu Christoff, 1997) with $2n = 14, 15$ and 16 (the latter diploid number has been cited as belonging to *Akodon* aff. *cursor* by Rieger *et al.*, 1995, and Geise *et al.*, 2001); *A. montensis* ($2n = 23$ - 26); *A. azarae* ($2n = 38$); *A. reinhardti* ($2n = 37,$

38); *A. lindberghi* ($2n = 42$); *A. serrensis* ($2n = 46$); *A. toba* ($2n = 42$ or 43) and *A. paranaensis* ($2n = 44$) (Yonenaga, 1972; Yonenaga, 1975; Yonenaga-Yassuda, 1979; Yonenaga-Yassuda *et al.*, 1987; Castro, 1989; Sbalqueiro, 1989; Geise *et al.*, 1996, 1998; Sbalqueiro and Nascimento, 1996; Fagundes *et al.*, 1998; Silva and Yonenaga-Yassuda, 1998; Christoff *et al.*, 2000).

Silva and Yonenaga-Yassuda (1998) suggested that the new $2n = 9, 10$ karyotype (*Akodon* sp. n.) from Mato Grosso state in central Brazil ranks as a new *Akodon* species. This karyotype has an autosomal polymorphism in chromosome 3, which can be either an acrocentric or a submetacentric due to a pericentric inversion, and the odd diploid number ($2n = 9$) being caused by monosomy of the X chromosome.

The $2n = 14, 15$ and 16 karyotypes have been assigned to the same taxonomic unit, *A. cursor* (Christoff, 1997; Fagundes *et al.*, 1998), but based on cytochrome-b mitochondrial gene sequences Geise *et al.* (2001) argued that $2n = 16$ specimens of this species from the Brazilian states of Paraná, Bahia and Minas Gerais should be recognized as being different from *A. cursor* and should be referred to as *Akodon* aff. *cursor*.

As the traditional classification of a large number of South American sigmodontine taxa remain poorly understood (Smith and Patton, 1999), chromosomal data and, more recently, mitochondrial DNA sequences (which al-

low better identification of species and phylogeographic patterns) have provided relevant information to establish relationships among this and other groups (Smith and Patton, 1993, 1999; Myers *et al.* 1995; Da Silva and Patton 1998; Garcia 1999; Geise *et al.* 2001; Bonvicino and Moreira 2001).

In this paper we compare G-band patterns between the $2n = 10$ and 16 *Akodon* karyotypes, and discuss the phylogenetic implications based on cytochrome-b sequences. Comparison of G-banded chromosomes, molecular systematics and biogeographic information provide additional data to aid our understanding of the evolutionary process as well as the present status and phylogeny of *Akodon*.

Material and Methods

Cytogenetic procedures

We used Seabright's method (1971) to obtain somatic cell G-banding patterns from 28 *Akodon* specimens with a $2n = 10$ karyotype and compared the G-banded chromosomes of $2n = 10$ specimens from Gaúcha do Norte in the Brazilian state of Mato Grosso and $2n = 16$ (referred to as *A. cursor* by Fagundes *et al.*, 1998) from the Brazilian state of São Paulo.

Sequencing procedures

Total genomic DNA from seven *Akodon* specimens with $2n = 10$ (specimens MZUSP-29658, 29671, 29673, 29678, 29684, 29685 and APC-270) was extracted from muscle or liver using the Chelex (BioRad) method (Walsh *et al.*, 1991). Tissues had previously been frozen in liquid nitrogen and transferred to 95% ethanol. The cytochrome-b gene was amplified by the polymerase chain reaction (PCR) with the MVZ05 and MVZ16 primer set as described by Smith and Patton (1993) and amplification confirmed by 2% agarose gel electrophoresis. The PCR products were sequenced using an ABI 377 automated sequencer using a d-Rhodamine cycle sequencing kit and the MVZ05 primer. The sequences were edited using the Sequence Navigator software (Applied Biosystems, Inc. 1994). Phylogenetic analyses under parsimony criterion were performed using the PAUP*4.0b6 program with 100 bootstrap replicates and Kimura-2-parameter distances (Kimura 1980) were also calculated.

The first 750 bp mitochondrial cytochrome-b sequences were obtained from seven *Akodon* specimens, four from Gaúcha do Norte and three from Vila Rica (both in Mato Grosso state), all specimens having been previously karyotyped. Since data indicated that these specimens share similar haplotypes we only use one specimen from each locality: the specimen MZUSP-29671 (GenBank accession number DQ631966) from Gaúcha do Norte and MZUSP-29685 (DQ631965) from Vila Rica in order to access the phylogenetic reconstruction. Other *Akodon* sequences and *Bolomys* (U03528) (*Necromys* after Smith and Patton,

1999) were obtained from GenBank, and the latter being used as the outgroup taxon. Voucher specimens with $2n = 9, 10$ were deposited at the Museu de Zoologia da Universidade de São Paulo (MZUSP), São Paulo state, Brazil.

Results and Discussion

Conventionally stained and banding patterns karyotypes of *Akodon* with $2n = 14, 15$ and 15 have been reported by various authors (Yonenaga, 1972, 1975; Yonenaga-Yassuda, 1979); Sbalqueiro and Nascimento, 1996; Fagundes *et al.*, 1998) and samples with a $2n = 9, 10$ karyotype have been described by Silva and Yonenaga-Yassuda (1998).

We compared G-bands of the $2n = 10$ and $2n = 16$ karyotypes, as fewer steps were involved in the differentiation between these karyotypes than between $2n = 10$ and $2n = 14$. This analysis suggests that a similar karyotype or a diploid number higher than $2n = 16$ gave rise to the lowest chromosome number of *Akodon* sp. n. ($2n = 10$). Diagrams of the $2n = 16$ and $2n = 10$ karyotypes are being shown in Figure 1.

Comparison of $2n = 10$ and 16 G-banded metaphases (Figure 2) revealed the following complex chromosomal rearrangements: chromosome 1 of $2n = 10$ could have derived from multiple tandem fusions of chromosomes 1 (after a pericentric inversion), 6 and 5q of a $2n = 16$ -like karyotype; chromosome 2 of $2n = 10$ could have originated by a Robertsonian rearrangement of chromosomes 3 (after a pericentric inversion) and 4 of $2n = 16$; chromosome 3 could have resulted from a tandem fusion involving chromosome 2 and probably 5p of $2n = 16$; pair 4 of $2n = 10$ and pair 7 of $2n = 16$ were the smallest chromosomes of each complement and seemed to be totally homologous in both karyotypes. The X chromosome was larger in the $2n = 10$




Figure 1 - Diagrammatic representation of *Akodon* karyotypes: (a) $2n = 10$ with pair 3 heteromorphic. (b) $2n = 16$.




Figure 2 - Comparison of G-banded karyotypes: *Akodon* with $2n = 10$ (left) and $2n = 16$ (right). Asterisks indicate centromere positions.

than in the $2n = 16$ karyotype due to the presence of a block of constitutive heterochromatin in the proximal region of the long arm and the Y chromosome was a minute subtelo-centric whereas in $2n = 16$ karyotype it was acrocentric.

Although we did not identify any homologies between the distal region of chromosome 3 of the $2n = 10$ karyotype and the other components of the $2n = 16$ karyotype it is possible that this sequence was lost from the ancestral $2n = 16$ karyotype to give rise to the $2n = 16$ karyotype and was then maintained in the $2n = 10$ karyotype, although the distal region of chromosome 3 could have arisen as a result of amplification of a segment in $2n = 10$ karyotype. Up to now we are not able to mention the origin, nature or function of this genetic material or if it implies difference in the fitness of the individuals. We can just mention that morphologically $2n = 10$ and $2n = 16$ seem to be indistinguishable (Christoff, in prep.)

Previous cytogenetic studies by Silva and Yonenaga-Yassuda (1998) revealed interstitial silver-stained nucleolar organizer regions (Ag-NORs) located at 1p and 1q, on the telomeric region of the 2p chromosome and the short arm of one pair 4 homologue of the $2n = 10$ karyotype, the $2n = 16$ karyotype reported by Fagundes *et al.* (1997) exhibited Ag-NORs on the telomeres of chromosome pairs 4 and 5. As far as we could detect, there was correspondence of Ag-NOR positions on the telomeric regions of both the 2p chromosome for the $2n = 10$ karyotype and the 4p chromosome for the $2n = 16$ karyotype. Absence of Ag-NORs on the 1p telomeres and their presence on one $2n = 10$ pair 4 homologue could be due to either loss of NORs or relocation of ribosomal genes.

According to Fagundes *et al.* (1997) fluorescence *in situ* hybridization analysis (FISH) with a telomeric (TTAGGG)_n probe resulted in only telomeric signals on the $2n = 16$ karyotype chromosomes. Interestingly, Silva and Yonenaga-Yassuda (1998) reported conspicuous inter-

stitial telomeric sites (ITS) on the pericentromeric region and the short arm of pair 1 and on the long arm of pair 3 of the $2n = 10$ karyotype, in addition to the regular telomeric signals. However these signals detected on the $2n = 10$ karyotype 1p and 3q chromosomes did not seem to be coincident with the specific positions involved in the chromosomal rearrangements. In addition, the pericentromeric signal of chromosome 1 was co-localized with pericentromeric constitutive heterochromatin, which is a highly repetitive DNA sequence. All this evidence led us to believe that much more complex rearrangements than we observe after the comparison of G-banded chromosomes had occurred during the evolutionary process which drove the differentiation of the $2n = 10$ and $2n = 16$ karyotypes.

Phylogenetic analyses and parsimony of cytochrome-b sequences support our previous chromosome results suggesting that the $2n = 10$ *Akodon* karyotype is a new species. These animals from Gaúcha do Norte and Vila Rica (localities geographically separated by nearly 450 km) share quite similar haplotypes (Kimura-2-parameter equal to 0.002) which are grouped in a strongly supported monophyletic group (bootstrap value of 100%), reinforcing the requirement of a new taxonomic status for this species (Figure 3). Furthermore, this species is recovered as the sister group (with bootstrap support of 100%) of *Akodon* specimens with $2n = 14, 15$ and 16 karyotypes, where the $2n = 16$ karyotype is the sister group of $2n = 14$ and 15 karyotypes. All of these three karyotypes show a common ancestor relative to other species of the *cursor*-complex, such as *A. montensis* and *A. mistax* (Figure 3).

Geise *et al.* (2001) reported a bootstrap value of 94% supporting *Akodon* with a karyotype of $2n = 14, 15$ as a sister of the $2n = 16$ karyotype, and referred to them as *A. cursor* and *Akodon* aff. *cursor*, respectively. After including our data ($2n = 10$ cytochrome-b sequences), the support for the relationships of these two taxa decreased to 72%, while




Figure 3 - Maximum parsimony majority consensus-tree (50% bootstrap) after a search with 100 replicates. Ninety-seven variable characters are parsimony uninformative and 558 are constant.

the bootstrap values to support the $2n = 14$, 15 and $2n = 16$ karyotypes as monophyletic groups increased respectively to 88 and 90%.

Kimura-2-parameter (K2p) distances within and among taxa were as follow: geographic samples of the $2n = 10$ karyotype differed by 0.002; $2n = 14$ - 15 by 0.007-0.023; $2n = 16$ by 0.032-0.052; and $2n = 10$ differed from $2n = 14$ - 15 by 0.053-0.062 and from $2n = 16$ by 0.057-0.071; $2n = 14$, 15 differed from $2n = 16$ by 0.040-0.067. *Akodon montensis* differed from them by 0.090-0.113 and *Necromys lasiurus* also differed from them by 0.155-0.190 K2p.

Maia and Langguth (1981) described an *Akodon* population from the Brazilian state of Pernambuco in which all specimens had a karyotype of $2n = 16$. More recently, *Akodon* specimens with a $2n = 16$ karyotype have also been found in the Brazilian states of Paraíba, Minas Gerais, São Paulo, Bahia and Paraná (Figure 4) (Rieger *et al.*, 1995; Sbalqueiro and Nascimento 1996; Fagundes *et al.*, 1998; Geise *et al.*, 2001). Regardless of the karyotype diversity in *Akodon* specimens with $2n = 14$, 15 and 16 karyotypes (a total of 28 karyotypes), Christoff (1997) and Fagundes *et al.* (1998), considered these specimens as a single species (*Akodon cursor*) and explained the variation as being due to pericentric inversions in three different chromosome pairs, a complex rearrangement involving centric fusion with previous pericentric inversions of two pairs, trisomy of pair 7 and monosomy of the X chromosome. But Christoff (1997) noted that *Akodon* specimens with $2n = 14$, 15 and 16 karyotypes were not distinctive taxonomically by external or cranio-dental morphology, and grouped as one unique species. However, the phylogenetic analyses of mitochondrial sequence data revealed, with relatively high support (72%), a node in which the *Akodon* specimens from São




Figure 4 - Distribution of *Akodon* with $2n = 10$; $2n = 14, 15$ and $2n = 16$, summarized from Yonenaga-Yassuda (1979); Maia and Langguth (1981); Rieger *et al.* (1995); Sbalqueiro and Nascimento (1996); Fagundes *et al.* (1998); Silva and Yonenaga-Yassuda (1998); Geise *et al.* (2001).

Paulo with a $2n = 14$ and 15 karyotype is the sister group of the *Akodon* specimens from Paraíba, Bahia and Minas Gerais with a karyotype of $2n = 16$. Moreover, according to Geise *et al.* (2001), cytochrome-b sequences of *Akodon* specimens from Una (Bahia state) with a $2n = 14$ - 15 karyotype share more similarities to specimens with the same karyotype which were 1500 km away than to $2n = 16$ karyotype specimens collected in the same locality. Therefore if the $2n = 14$ and 16 karyotypes were panmictic (randomly interbreeding) there should be no reason for their mitochondrial DNA to appear so different as the results show.

Rieger *et al.* (1995) used 22 structural loci encoding 14 proteins to check genetic variability in 22 *Akodon* specimens from Paraná with $2n = 16$ karyotypes, 41 specimens from Espírito Santo with $2n = 14$ and 15 karyotypes and 206 specimens from the Brazilian states of Santa Catarina and three localities in the Brazilian state of Rio Grande do Sul with $2n = 24$, 25 and 26 karyotypes. On the basis of phenetic and phylogenetic data, Rieger *et al.* (1995) suggested that members of the *cursor*-complex have undergone recent differentiation and that chromosomal divergence has proceeded without eletromorphic divergence. We can add here that mitochondrial sequences can identify and separate these three units plus the new species with the $2n = 10$ karyotype.

It is also interesting to point out that preliminary morphometric studies (Christoff *et al.*, in prep.) showed no differences between $2n = 10$ karyotype *Akodon* and *A. cur-*

sor (sensu Christoff, 1997), indicating the possibility that they are cryptic species as already observed in *A. cursor* and *A. montensis* (Christoff, 1997).

It therefore seems that a special evolutionary situation exists in which a group is recognized as a unique species on the basis of morphological traits but karyologically is clearly separated into $2n = 14$, 15 and 16 forms and $2n = 10$, with cytochrome-b sequences indicating that the $2n = 10$ karyotypes represents a well supported full species and $2n = 24$ is another strongly bootstrapped species while *Akodon* with 16 chromosomes is probably a different entity distinct from *Akodon* with $2n = 14$ - 15 chromosomes as previously hypothesized by Geise *et al.* (2001).

Acknowledgments

We thank Dr. Yuri Leite for the valuable contributions during the lab-work, data analysis, and elaboration of the manuscript. We are also grateful to Dr. Valéria Fagundes for the concession of $2n = 16$ G-bands, Dr. Katia Pellegrino and Sheda Morshed for suggestions in the manuscript; MSc. Ana Paula Carmignotto and Dr. Alexandre U. Christoff who collected specimens, and to Dr. Miguel T. Rodrigues, Dr. Vinícius Xavier da Silva, Dr. Gabriel Skuk and MSc. Dante Pavan for helping in fieldwork; to Miriam Romeo, Tais Machado and Yukie Sato for technical assistance. We also thank Dr. Yuri Leite, Margareth Smith and Lena Geise for providing sequences which are also available in the GenBank, and the MVZ for the accessibility and infra-structure. Grants from Fundação de Amparo à Pesquisa do Estado de São Paulo (including 00/06591-0 to Dr. M.J.J.S. and BIOTA 98/5075 to Dr. Mario de Vivo); Coordenação de Aperfeiçoamento de Pessoal de Nível Superior, Conselho Nacional de Desenvolvimento Científico e Tecnológico and National Science Foundation (grants to Dr. James Patton) are also acknowledged.

References

- Bonvicino CR and Moreira MAM (2001) Molecular phylogeny of the genus *Oryzomys* (Rodentia, Sigmodontinae) based on cytochrome-b DNA sequences. *Mol Phylogenet Evol* 18:282-292.
- Castro EC (1989) Ocorrência e caracterização cromossômica de roedores Akodontinos do Rio Grande do Sul. Dissertação de Mestrado, Universidade Federal do Rio Grande do Sul, Porto Alegre.
- Christoff AU, Fagundes V, Sbalqueiro IJ, Mattevi MS and Y Yonenaga-Yassuda Y (2000) Description of a new species of *Akodon* (Rodentia, Sigmodontinae) from Southern Brazil. *J Mammalogy* 81:838-851.
- Christoff AU (1997) Contribuição à sistemática das espécies do gênero *Akodon* (Rodentia, Sigmodontinae) do leste do Brasil: Estudos anatômicos, citogenéticos e de distribuição geográfica. Tese de Doutorado, Instituto de Biociências, Universidade de São Paulo, São Paulo.
- Da Silva MNF and Patton JL (1998) Molecular phylogeography and the evolution and conservation of Amazonian mammals. *Mol Ecol* 7:475-486.
- Fagundes V and Yonenaga-Yassuda Y (1998) Evolutionary conservation of the whole homologous chromosome arms in the Akodont rodents *Bolomys* and *Akodon* (Muridae, Sigmodontinae): Maintenance of interstitial telomeric segments (ITBs) in recent event of centric fusion. *Chromosome Res* 6:643-648.
- Fagundes V, Christoff AU and Yonenaga-Yassuda Y (1998) Extraordinary chromosomal polymorphism with 28 different karyotypes in the neotropical species of *Akodon cursor* (Muridae, Sigmodontinae), one of the smallest diploid numbers in rodents ($2n = 16$, 15 and 14). *Hereditas* 129:263-274.
- Fagundes V, Vianna-Morgante AM and Yonenaga-Yassuda Y (1997) Telomeric sequences localization and G-banding patterns in the identification of a polymorphic chromosomal rearrangement in the rodent *Akodon cursor* ($2n = 14$, 15 and 16). *Chromosome Res* 5:228-232.
- Garcia LF (1999) Molecular phylogenetics of neotropical oryzomyine rodents (Muridae, Sigmodontinae). Ph.D. Thesis, University of California, Berkeley.
- Geise L, Canavez FC and Seuánez HN (1998) Comparative karyology in *Akodon* (Rodentia, Sigmodontinae) from Southwestern Brazil. *J Hered* 89:158-163.
- Geise L, Cerqueira R and Seuánez HN (1996) Karyological characterization of a new population of *Akodon lindberghi* (Rodentia, Sigmodontinae) in Minas Gerais state (Brazil). *Caryologia* 49:57-63.
- Geise L, Smith MF and Patton JL (2001) Diversification in the genus *Akodon* (Rodentia, Sigmodontinae) in southeastern America: Mitochondrial DNA sequences analysis. *J Mammalogy* 82:92-101.
- Kasahara S and Yonenaga-Yassuda Y (1984) A progress report of cytogenetic data on Brazilian rodents. *Rev Bras Genét* 7:509-533.
- Kimura M (1980) A simple method for estimating evolutionary rate of base substitution through comparative studies of nucleotide sequences. *J Mol Evol* 16:111-120.
- Liascovich RC and Reig OA (1989) Low chromosomal number in *Akodon cursor montensis* Thomas, and karyologic confirmation of *Akodon serrensis* Thomas in Misiones, Argentina. *J Mammalogy* 70:391-395.
- Maia V and Langguth A (1981) New karyotypes of Brazilian Akodont with notes on taxonomy. *Zeitschr Säugetierkunde* 46:241-249.
- Myers P, Lundrigan B and Tucker PK (1995) Molecular phylogenetics of oryzomyine rodents: The genus *Oligoryzomys*. *Mol Phylogenet Evol* 4:372-382.
- Rieger TT, Langguth A and Wienmer TA (1995) Allozymic characterization and evolutionary relationships in the Brazilian *Akodon cursor* species group (Rodentia, Cricetidae). *Biochem Genet* 33:283-295.
- Sbalqueiro IJ and Nascimento AP (1996) Occurrence of *Akodon cursor* (Rodentia, Cricetidae) with 14 , 15 and 16 chromosome karyotypes in the same geographic area in South Brazil. *Braz J Genet* 19:565-569.
- Seabright M (1971) A rapid banding technique for human chromosomes. *Lancet* 2:971-972.
- Silva MJJ and Yonenaga-Yassuda Y (1998) Karyotype and chromosomal polymorphism of an undescribed *Akodon*

- from Central Brazil, a species with the lowest diploid chromosome number in rodents. *Cytogenet Cell Genet* 81:46-50.
- Smith MF and Patton JL (1999) Phylogenetic relationships and the radiation of sigmodontine rodents in South America: Evidence from cytochrome b. *J Mammal Evol* 6:89-128.
- Smith MF and Patton JL (1993) The diversification of South American murid rodents: Evidence from mitochondrial DNA sequence data for the akodontine tribe. *Biol J Linnean Soc* 50:149-177.
- Spotorno OA and Fernandez R (1976) Chromosome stability in southern *Akodon* (Rodentia, Cricetidae). *Mammalian Chromosome Newsletter* 17:13-14.
- Svartman M and Almeida EJC (1994) The karyotype of *Akodon lindberghi* Hershkovitz, 1990 (Cricetidae, Rodentia) from Central Brazil. *Rev Bras Genét* 17:963-972.
- Walsh PS, Metzger DA and Higuchi R (1991) Chelex 100 as a medium for simple extraction of DNA for PCR-based typing from forensic material. *Biotechniques* 10:506-513.
- Yonenaga Y (1972) Chromosomal polymorphism in the rodent *Akodon arviculoides* ssp. ($2n = 14$) resulting from two pericentric inversions. *Cytogenetics* 11:448-499.
- Yonenaga Y (1975) Karyotypes and chromosome polymorphisms in Brazilian rodents. *Caryologia* 28:269-286.
- Yonenaga-Yassuda Y (1979) New karyotypes and somatic and germ-cell banding in *Akodon arviculoides* (Rodentia, Cricetidae). *Cytogenet Cell Genet* 23:241-249.
- Yonenaga Y, Frota-Pessoa O, Kasahara S and Almeida EJC (1976) Cytogenetic studies on Brazilian rodents. *Ciência & Cultura* 28:202-211.
- Yonenaga-Yassuda Y, Pereira LA, Armada JL and L'Abbate M (1987) Chromosomal polymorphism in *Akodon reinhardti* Langguth, 1975 (Rodentia, Cricetidae). *Rev Bras Genét* 10:199-208.

Associate Editor: João S. Morgante