

Autosomal and sex chromosomal polymorphisms with multiple rearrangements and a new karyotype in the genus *Rhipidomys* (Sigmodontinae, Rodentia)

MARIA JOSÉ J. SILVA and Y. YONENAGA-YASSUDA

Departamento de Biologia, Instituto de Biociências, Universidade de São Paulo, SP, Brazil

Silva, M. J. J. and Yonenaga-Yassuda, Y. 1999. Autosomal and sex chromosomal polymorphisms with multiple rearrangements and a new karyotype in the genus *Rhipidomys* (Sigmodontinae, Rodentia).—*Hereditas* 131: 211–220. Lund, Sweden. ISSN 0018-0661. Received December 9, 1998. Accepted November 11, 1999

Two diploid numbers and five karyomorphs were found in ten specimens of *Rhipidomys* (Sigmodontinae, Rodentia) from three states in Brazil: $2n = 50$ from Amazonas, and $2n = 44$ from Mato Grosso and Bahia. CBG, GTG, and RBG-banding and Ag-NOR analyses were performed, as well as fluorescence in situ hybridization with $(T_2AG_3)_7$ probes. The new diploid number of $2n = 50$ was associated with two different fundamental numbers (FN = 71 and 72) as a result of pericentric inversions and addition/deletion of constitutive heterochromatin. The samples from two localities (Aripuanã and Vila Rica) in the state of Mato Grosso shared $2n = 44$ and FN = 52, but their karyotypes differed because of pericentric inversions. Although the single specimen from Bahia had the same diploid number as the samples from Mato Grosso, its karyotype and FN were completely distinctive. Karyological comparison of GTG-banding patterns revealed total homology between the karyotypes of the specimens from Bahia and Mato Grosso, implying the occurrence of 14 autosomal pericentric inversions. Homologies between ten of the autosomes in the karyotypes with $2n = 50$ (FN = 72) and $2n = 44$ (from Vila Rica, MT) were demonstrated. The differentiation between $2n = 44$ and $2n = 50$ involved five pericentric inversions, addition/deletion of constitutive heterochromatin in both autosomes and sex chromosomes, at least one Robertsonian rearrangement and other not detected rearrangements. Despite the remarkable number of rearrangements, interstitial telomeric sites (ITS) were not detected. Sex chromosomes also exhibited polymorphism in size and morphology.

Yatiyo Yassuda, Departamento de Biologia, Instituto de Biociências, Universidade de São Paulo, São Paulo, São Paulo – Brazil – 05422-970, C.P. 11.461. E-mail: mariajo@usp.br

Climbing rats, genus *Rhipidomys* (Sigmodontinae, Rodentia), occur in forest, gardens and plantations and also invade rural camps and houses throughout South America and Panama (EMMONS and FEER 1997). According to CABRERA (1961), the genus comprised five species, four of which were polytypic; HONACKI et al. (1982) and NOWAK (1991) recognized seven species; REIG (1986) identified nine species; MUSSEY and CARLETON (1993) 14, and TRIBE (1996) recognized 18 species. No recent critical synopses of valid species have been published and a broad systematic revision is needed.

Four of the 14 species accepted by MUSSEY and CARLETON (1993) are found in Brazil: *Rhipidomys leucodactylus* (synonyms: *aratayae*, *bovallii*, *equatoris*, *lucullus*, *rex*, *sclateri*) in the north; *R. mastacalis* (synonyms: *cearanus*, *emiliae*, *macrurus*, *maculipes*, *yuruanus*) in central and eastern Brazil; *R. nitela* (synonyms: *fervidus*, *milleri*, *tobagi*) in north-central Brazil; and *R. macconnelli* (synonym: *subnubis*) perhaps in the northernmost region of the country. On the other hand, TRIBE (1996) considered that there are nine species and one *incertae sedis* in Brazilian territory: *R. leucodactylus*, *R. macconnelli*, *R. nitela*, *R. emiliae*, *R. macrurus*, *R. mastacalis*, *R. cf. macrurus*, *R. sp. 4*, *R. sp. 2*, *R. inc. sedis*.

Heretofore, cytogenetic studies of *Rhipidomys* specimens performed by GARDNER and PATTON (1976), ZANCHIN et al. (1992), SVARTMAN and ALMEIDA (1993), and CORRÊA and PESSÔA (1996) revealed $2n = 44$ as the common diploid number found in one locality from Colombia and eight sites from Brazil (Fig. 1). Although the chromosome number was constant, variation occurred in the fundamental number (FN).

Herein we present five karyomorphs (two from Amazonas, one from Bahia, and two from Mato Grosso), including CBG, GTG, and RBG-banding, and Ag-NOR analysis in ten specimens of *Rhipidomys*, as well as fluorescent in situ hybridization (FISH) with a telomeric probe.

MATERIALS AND METHODS

Specimens

Specimens identified preliminarily during fieldwork as *Rhipidomys* cf. *mastacalis* were captured in the Amazonian rain forest of the state of Mato Grosso: two males and four females at Vila Rica (9°54'28"S, 51°12'13"W) and one female at Aripuanã (10°10'S, 59°27'W). One female of *Rhipidomys* sp. A was

trapped at Morro do Chapéu (11°36'S, 41°08'W), state of Bahia at an altitude of about 1000 m in the rocky habitats called "campos rupestres". Two males of *Rhipidomys* sp. B were collected in the Amazonian rain forest near Manaus (2°25'S, 59°50'W), state of Amazonas. Fig. 1 includes these data.

Skulls and skins of the specimens from Mato Grosso and Bahia were deposited in the collection of the Museu de Zoologia, Universidade de São Paulo (MZUSP), Brazil.

Chromosome preparation and banding

Chromosome preparations were obtained from bone marrow and spleen after in vivo colchicine treatment. Fibroblast cultures from ear and tail biopsies from five individuals were established in the laboratory using Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 20% fetal bovine serum. CBG, GTG, and RBG-banding patterns, as well as Ag-NOR staining, were carried out using routine cytogenetic techniques, according to SUMNER (1972), SEABRIGHT (1971), DUTRILLAUX and COUTURIER (1981), and HOWELL and BLACK (1980), respectively.

Fluorescence in situ hybridization (FISH)

Oligonucleotides (T₂AG₃)₇ in both sense and anti-sense orientation (GIBCO BRL) were used as probes for FISH, after 3'-end labeling with biotin-11-dUTP (Boehringer) for localization of telomeric sequences. Hybridization was performed overnight at 37°C and the signals were detected by incubation with fluores-

cein isothiocyanate (FITC)-labeled anti-biotin. FISH was also performed with the commercially available digoxigenin labeled "all human telomeres probe" (P5097-DG5, Oncor). The slides were counterstained with propidium iodide and mounted with an antifade solution. A total of 40 metaphases from six specimens was analyzed using a Zeiss Axiophot microscope equipped with a FITC filter and photographed with Ektachrome 400 (Kodak).

RESULTS

Rhipidomys cf. *mastacalis*

Rhipidomys cf. *mastacalis* from the two localities in the state of Mato Grosso presented a karyotype with $2n = 44$, $FN = 52$ (Fig. 2a) with 16 uniarmed and 5 biarmed pairs of autosomes. Despite the similarities in diploid and fundamental numbers, two different karyotypes were detected on the basis of variations in the biarmed pairs. In the sample from Vila Rica (Fig. 2a) pair 4 and three small pairs (19 to 21) were biarmed, while in the Aripuanã specimen (Fig. 3a) pair 4 was uniarmed and four small pairs were biarmed (18 to 21). Pair 10 was biarmed in both karyomorphs. The X chromosome was acrocentric (Xa); the Y chromosome was acrocentric (Ya) and was the same size as pair 21.

The CBG-banding pattern revealed very small amounts of constitutive heterochromatin in the pericentromeric region of autosomes (Fig. 3a). The Xa




Fig. 1. Localities, diploid and fundamental numbers of *Rhipidomys*. ■ GARDNER and PATTON (1976); □ ZANCHIN et al. (1992); ◆ SVARTMAN and ALMEIDA (1993), ● CORRÊA and PESSÔA (1996) ▲ Present work. (*) Although the fundamental number is the same ($FN = 52$) in the specimens from MT, the biarmed homologues are different in the two karyotypes.




Fig. 2. a–d. Conventionally stained karyotypes of *Rhipidomys* (bar = 10 μ m). **a** Male of *R. cf. mastacalis* from Vila Rica, $2n = 44$ (FN = 52) and XaYa. Inset: XaXa. **b** Female of *R. sp. A*, $2n = 44$ (FN = 61) with nine heteromorphic pairs and XaXb. **c** Male of *R. sp. B*, $2n = 50$ (FN = 71, heteromorphic for n°8) and XcYb. **d** Male of *R. sp. B*, $2n = 50$ (FN = 72) and XcYc.

and Ya chromosomes also exhibited small pericentromeric C-bands. RBG-banding revealed the late replicating X chromosome and allowed precise identification of Xa (Fig. 4a).

The Ag-NOR analysis of 60 metaphases from the Vila Rica sample revealed NORs in the short arms of one large pair (no. 3) and three small pairs—probably pairs 12, 15 and 19 (Fig. 5a). The NORs ranged from 4 to 8 per cell, with a mean and mode of 7 NORs. In the specimen from Aripuanã, 11 metaphases showed variation from 7 to 12 Ag-NORs, with a mean and mode of 8 NORs, located in the short arms of one large pair and the short arms of some small autosomes. Additionally, one homologue of the smallest

chromosome, the biarmed pair 21, was Ag-stained (Fig. 5a). Associations involving NOR-bearing chromosomes were observed in both kinds of karyotypes.

Rhipidomys sp. A

Rhipidomys sp. A from Bahia had $2n = 44$ and FN = 61. The karyotype comprised seven heteromorphic pairs (1 to 5, 8 and 13) composed of one biarmed and one unarmed homologue; pair 13 was also heteromorphic in size; pairs 7 and 9 were heteromorphic with submetacentric and submetacentric homologues in the former and submetacentric and metacentric homologues in the latter (Fig. 2b). Pairs 6 and 10 to 12 were biarmed. Pair 14, the largest autosome pair

in the complement, and pairs 15 to 21 were acrocentric. The X chromosomes were heteromorphic: one was acrocentric (Xa) and the other was submetacentric (Xb). Xa had a very small short arm, which distinguished it from the autosomal acrocentrics. All autosomes could be identified not by conventional staining but based on banding patterns.

CBG-banding showed that constitutive heterochromatin was almost absent in the largest pairs but was present in the pericentromeric regions of certain small autosomes (Fig. 3b). Xa and Xb had conspicuous blocks in their pericentromeric regions. The heterochromatin in the short arm of Xb stained less than the pericentromeric heterochromatin. RBG and GTG-banding allowed identification of all the autosomal pairs. Incorporation of BrdU in fibroblast cultures showed that Xb was inactivated in all the 22 metaphases analyzed (Fig. 4b).

From a total of 35 metaphases, 29 presented five Ag-NORs and 6 showed four AgNORs, which were located at the end of one homologue of pair 9, in the long arm of the acrocentric homologue of pair 13, and in the short arms of three medium-sized acrocentric pairs, probably nos. 16 and 18 (Fig. 5b).




Fig. 3. a–c. CBG banded karyotypes of *Rhipidomys*. a Female of *R. cf. mastacalis* from Aripuanã, $2n = 44$ (FN = 52), XaXa. Inset: XaYa. b Female of *R. sp. A*, $2n = 44$ (FN = 61) and XaXb. c Male of *R. sp. B*, $2n = 50$ (FN = 71) and XcYc. Inset: XcYc.




Fig. 4. a–c. Partial metaphases of *Rhipidomys* after RBG-banding. **a** Female of *R. cf. mastacalis* from Vila Rica, $2n = 44$ (FN = 52), XaXa. Asterisk indicates the late replicating Xa. **b** Female of *R. sp. A* with $2n = 44$ (FN = 61) showing the late replicating Xb. **c** Male of *R. sp. B* with $2n = 50$ (FN = 71) exhibiting late replicating Yb and short arm of Xc.

Rhipidomys sp. B

Two males of *Rhipidomys sp. B* from Manaus had $2n = 50$, but two different fundamental numbers, FN = 71 and 72 (Fig. 2c, d). The specimen with FN = 71 had 11 biarmed pairs (1 to 7 and 9 to 12), 12 uniarmed pairs (13 to 24) and a heteromorphic pair 8 (one biarmed and one uniarmed). The specimen with FN = 72 showed 12 biarmed pairs (1 to 12) and 12 uniarmed pairs of autosomes (13 to 24). The X chromosome was a large submetacentric (Xc) in both individuals, and the Y displayed two morphologies: a small acrocentric (Yb) in the specimen with FN = 71 and a medium-sized acrocentric (Yc) with a very small short arm in the other individual.

CBG-banding revealed the presence of weakly stained heterochromatin in the pericentromeric region of some autosomes and in the short arm of one homologue of the heteromorphic pair 8 in the specimen with FN = 71 (Fig. 3c). This pair is probably not equivalent to pair 8 in the specimen with FN = 72, which did not exhibit heterochromatin in its short arm. The short arm of Xc was heterochromatic and its long arm had two C-positive interstitial bands; Yb was heterochromatic in its pericentromeric region and the distal part of its long arm; and Yc was entirely heterochromatic (Fig. 3c). RBG-banding revealed late replication of the Y and of the short arm of the X chromosome (Fig. 4c).

Ag-NORs were analyzed in 50 metaphases of the specimen with FN = 71 and ranged from 4 to 6, with a mean and mode of 5 NORs. The NOR-bearing chromosomes were one homologue of pair 5 and one

of 16, and both homologues of pairs 22 and 24. Associations were observed (Fig. 5c). In the specimen with FN = 72, analysis of 27 metaphases evinced from 3 to 6 Ag-NORs, with a mean and mode of 4 NORs. The NOR bearing chromosomes were one homologue of pair 5, one of pair 16, and both homologues of pairs 10 and 24 (Fig. 5d). Pair 5 exhibited one secondary constriction in the short arm, in which the NORs were located. Associations were not detected. Comparison of the Ag-NOR-bearing chromosomes in the FN = 71 and 72 karyotypes revealed that the difference between them was due to the presence of ribosomal cistrons in the short arms of pairs 10 and 22, respectively.

All these data together led us to suppose that pairs 8, 9, 10 and 11 could not correspond to the homologous with the same numbers in both karyotypes (FN = 71 and 72). Probably pair 8 from the karyotype with FN = 71 belongs to the acrocentric group in the FN = 72, and pair 10, which is a metacentric in the FN = 71, is different from the subtelocentric pair 10 of FN = 72 karyotype (Fig. 2c and 2d).

Comparative GTG-banding pattern

A detailed karyological comparison of GTG-banded chromosomes revealed total homology between the $2n = 44$ karyotypes of *Rhipidomys cf. mastacalis* from Vila Rica and *R. sp. A* from Bahia, implying that their differences were the result of pericentric inversions in 14 autosome pairs (Fig. 6).

Comparison between the karyotypes of *R. cf. mastacalis* and *R. sp. B* with $2n = 50$, FN = 72 (Fig. 6) showed the homology of 10 pairs of autosomes and

revealed five pericentric inversion events. Chromosome pairs 7, 9, 11, and 14 to 24 of the $2n = 50$ karyotype could not be identified in the $2n = 44$ karyotypes. The X chromosomes presented two characteristic dark bands in their long arms after G-banding. The Yb showed two dark bands, one pericentromeric and one distal, after G-banding and the Yc had the same pattern and also a positively stained short arm.

FISH

Hybridization signals were observed in the telomeres of all chromosomes of *Rhipidomys* cf. *mastacalis*, *R.* sp. A and *R.* sp. B. No interstitial telomeric sites (ITS) were seen (Fig. 7).

DISCUSSION

Cytogenetic and taxonomic research on the genus *Rhipidomys* is almost absent in the literature and few cytogenetic studies have been performed on Brazilian populations. Karyological information is available for around 21 specimens, ten of which are described herein. Our studies reveal a high level of polymorphism with five different karyomorphs, including two karyotypes with $2n = 50$ and three with $2n = 44$.

The karyotypes of *R.* sp. B represent a previously unreported diploid number for the genus, and the difference between the fundamental numbers found in this species is due to two different chromosomal rearrangements: pericentric inversion and addition/




Fig. 5. a–d. Ag-NO₃ stained karyotypes of *Rhipidomys*. a *R.* cf. *mastacalis* from Vila Rica ($2n = 44$, FN = 52) with 6 Ag-NORs. Inset: pair 21 of the specimen from Aripuanã. b *R.* sp. A ($2n = 44$, FN = 61) with 5 Ag-NORs. c *R.* sp. B ($2n = 50$, FN = 71) with 6 Ag-NORs. d *R.* sp. B ($2n = 50$, FN = 72) with 5 Ag-NORs. Arrows indicate associations involving pairs 3 and 12 in (a) and pairs 22 and 24 in (c).




Fig. 6. Comparison of GTG-banded haploid sets of *Rhipidomys*. From left to right in each triplet: (⊕) *R* sp. B ($2n = 50$, FN = 72); (⊖) *R. cf. mastacalis* from Aripuanã ($2n = 44$, FN = 52); (⊕) *R. sp. A* ($2n = 44$, FN = 61). Inset: Yc and Yb of *R. sp. B*.

deletion of constitutive heterochromatin. The former mechanism probably involves pair 10 from the FN = 72 karyotype and its counterpart, pair 22, from the FN = 71 complement. This hypothesis could be corroborated by the observation that the NOR-bearing chromosomes are the same in both karyotypes, except for pairs 10 and 22. It led us to hypothesize that pair 10 of the FN = 71 karyotype is equivalent to pair 22 of the FN = 72 karyotype, and that a pericentric inversion would be involved in the differentiation of this pair.

The other mechanism responsible for the variability in fundamental numbers is addition/deletion of constitutive heterochromatin in the short arm of one homologue of pair 8 from the specimen with FN = 71. Since there is no similar heterochromatic pattern in the FN = 72 specimen, this pair probably corresponds to an acrocentric pair in the latter karyotype.

Fundamental number variability in *Rhipidomys cf. mastacalis* resulted from pericentric inversions: the six individuals from Vila Rica presented a biarmed homomorphic pair 4, whereas this pair was homomorphic and uniarmed in the specimen from Aripuanã; a

heteromorphic pair 4 was not observed in our sample. A pericentric inversion involving one small pair in the specimen from Aripuanã was also detected and resulted in a karyotype with four small biarmed pairs, whereas the karyotype of the sample from Vila Rica included only three small biarmed pairs. Pair 10 was biarmed in all these specimens from the state of Mato Grosso. A heteromorphism of pair 10 in individuals from São Paulo and Brasília was reported by SVARTMAN and ALMEIDA (1993).

Variability in the number of Ag-NORs between the Vila Rica and Aripuanã samples could reflect either a real difference in the number of nucleolar organizer regions or differential activity of ribosomal cistrons. This could also be the case with the smallest stained metacentric in pair 21 of the Aripuanã specimen.

Among the specimens with 44 chromosomes, *Rhipidomys sp. A* presented an unexpected and exceptional karyotype with nine heteromorphic autosomal pairs and heteromorphic X chromosomes. Considering the high number of heteromorphic pairs, we hypothesize that: (1) the *Rhipidomys sp. A* individual could be from a population in which these hetero-

The X and Y chromosomes in our sample were variable in size and morphology (Xa, Xb and Xc and Ya, Yb and Yc) because of additions/deletions of constitutive heterochromatin. Our Xa, Xb and Ya were similar, respectively, to the Xa, Xb and Y described by SVARTMAN and ALMEIDA (1993). However, the short arm of our Xc was larger than the short arm of Xb + described by the same authors.

All specimens of *Rhipidomys* for which karyotypes have so far been published (Fig. 1) have had the same diploid number of $2n = 44$, with variable fundamental numbers due to the occurrence of pericentric inversions. Although skull and skin data show a clear difference between the samples, the karyotype of *R. leucodactylus* from Rondônia (FN = 48; uniarmed pair 4 and 10) (ZANCHIN et al. 1992), is similar to that of a female *Rhipidomys* sp. from Brasília (FN = 48) (SVARTMAN and ALMEIDA 1993) and a specimen from Mocambinho (FN = 48) (CORRÊA and PESSÔA 1996), except in its sex chromosome morphology. A *Rhipidomys* sp. from São Paulo has an autosomal set similar to that of a male from Brasília ($2n = 44$, FN = 49 and pair 10 heteromorphic) (SVARTMAN and ALMEIDA 1993). These karyotypes differ from that of *Rhipidomys* cf. *mastacalis* from Vila Rica only in the morphology of pairs 4 and 10, which are biarmed in the latter species. All the karyotypes mentioned above differ from the complement found in a specimen from the state of Espírito Santo (ZANCHIN et al. 1992), which presented four biarmed chromosome pairs, and from the karyotype of the Aripuanã specimen, which had five biarmed pairs.

TRIBE (1996) identified the *Rhipidomys* specimens from São Paulo and Espírito Santo as *R.* cf. *macrurus*, and those from Brasília and Mocambinho as *R. macrurus*. He considered our *R.* sp. B preliminarily to be *R. nitela*. A detailed systematic study of the different forms of *Rhipidomys*, including our samples, seems urgent.

Our cytogenetic studies suggest that our *R.* sp. B represents a distinctive species, separate from *R.* cf. *mastacalis* and *R.* sp. A. Moreover, there are two sympatric species in Mato Grosso state: the smaller *R. nitela* (MUSSEY and CARLETON 1993; TRIBE 1996) and the larger *R. emiliae* (TRIBE 1996) or *R. mastacalis* (MUSSEY and CARLETON 1993). Morphological studies should be performed on our samples preliminarily identified as *R.* cf. *mastacalis* in order to clarify this point. More *Rhipidomys* specimens should also be studied from the "campos rupestres" of Bahia, a habitat different from the Atlantic and Amazonian rain forests, in order to ascertain whether our *R.* sp. A might represent a hybrid in the genus.

ACKNOWLEDGEMENTS

We are indebted to Ana Paula Carmignotto, Dr. Pedro Luís Bernardo da Rocha and Dr. Jay R. Malcolm, who collected the specimens; to Dr. Marta Svartman and Eunice J. Cardoso de Almeida and especially to Dr. Christopher Tribe for critical reading of the manuscript; to Tien Hsi Chu, Miriam Romeo and Tais Machado for technical assistance; to Seplan (Secretaria de Planejamento do Estado de Mato Grosso) and CNEC (Consórcio Nacional de Engenheiros Consultores) for supporting the fieldwork in the state of Mato Grosso. The specimens from Manaus were collected during the Project "Dinâmica Biológica de Fragmentos Florestais" supported by INPA (Instituto Nacional de Pesquisas da Amazônia) and WWF (World Wildlife Foundation) under Dr. Jay Malcolm's supervision. We are especially grateful to Dr. Miguel T. Rodrigues and his team for helping in the fieldwork and to Gabriel Skuk for helping in the elaboration of the map. This work was also supported by grants from the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), Financiadora de Estudos e Projetos (FINEP), Fundação Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq).

REFERENCES

- Cabrera A, (1961). Catálogo de los mamíferos de America del Sur. Rev. Mus Argent. Cienc. Nat. 6: 309–732.
- Corrêa MMO and Pessôa LM, (1996). O cariótipo de *Rhipidomys* sp. (Rodentia: Cricetidae) do norte de Minas Gerais. Resumos do XXI Congresso Brasileiro de Zoologia, Porto Alegre, 5–9 fey, 1061, p. 226.
- Dutrillaux B and Couturier J, (1981). La pratique de l'analyse chromosomique. Mason. Paris.
- Emmons LE and Feer F, (1997). Neotropical rainforest mammals. A field guide. 2nd edition. University of Chicago. Chicago.
- Garagna S, Broccoli D, Redi CA, Scarle JB, Cooke HJ and Capanna E, (1995). Robertsonian metacentrics of house mouse lose telomeric sequences but retain some minor satellite DNA in the pericentromeric area. Chromosoma 103: 685–692.
- Gardner AL and Patton JL, (1976). Karyotypic variation in *Oryzomyine* rodents (Cricetidae) with comments on chromosomal evolution in the Neotropical Cricetine Complex. Occ. Pap. Mus Zool. LA State University 49: 1–48.
- Honacki JH, Kinman KE and Koepl JW, (1982). Mammals species of the world. Allen Inc. and the Association of Systematics Collections. Lawrence, Kansas, USA.
- Howell WM and Black DA, (1980). Controlled silver-staining of nucleous organizer regions with a protective colloidal developer: a 1-step method. Experientia 36: 1014–1015.
- Musser GG and Carleton MD, (1993). Family Muridae. In: Mammal Species of the World: a Taxonomic and Geographic Reference. (eds DE Wilson and DM Reeder). 2nd ed. Smithsonian Institution. Washington/London.
- Nanda I, Schneider-Rasp S, Winking H and Schmid M, (1995). Loss of telomeric sites in the chromosomes of *Mus musculus domesticus* (Rodentia, Muridae) during Robertsonian rearrangements. Chrom. Res. 3: 399–409.

- Nowak RM, (1991). Walker's Mammals of the World. 5th ed., vol. II, 1629 p. John Hopkins University. Baltimore and London.
- Reig AO, (1986). Diversity patterns and differentiation of high Andean rodents. In: High altitude tropical biogeography, (eds F Vuilleumier and Monasterio) New York. Oxford Univ.: 404–439.
- Seabright M, (1971). A rapid banding technique for human chromosomes. *Lancet* 2: 971–972.
- Silva MJJ and Yonenaga-Yassuda Y, (1998). Heterogeneity and meiotic behaviour of B and sex chromosomes, banding patterns and localization of (TTAGGG)_n sequences by FISH, in the Neotropical water rat *Nectomys* (Rodentia, Cricetidae). *Chrom. Res.* 6: 455–462.
- Silva MJJ and Yonenaga-Yassuda Y, (1997). New karyotypes of two related species of *Oligoryzomys* genus (Cricetidae, Rodentia) involving centric fusion with loss of NORs and distribution telomeric (TTAGGG)_n sequences. *Hereditas* 127: 217–229.
- Sumner AT, (1972). A simple technique for demonstrating centromeric heterocromatin. *Exp Cell Res.* 75: 304–306.
- Svartman M and Almeida EJC, (1993). Pericentric inversion and X chromosome polymorphism in *Rhipidomys* sp. (Cricetidae, Rodentia) from Brazil. *Caryologia* 46: 219–225.
- Tribe CJ, (1996). The Neotropical rodent genus *Rhipidomys* (Cricetidae, Sigmodontinae) – a taxonomic revision. PhD dissertation. University College London.
- Zanchin NIT, Langguth A and Mattevi MS, (1992). Karyotypes of Brazilian species of *Rhipidomys* (Rodentia, Cricetidae). *J Mamm.* 73: 120–122.