

A NEW RECORD AND OBSERVATIONS OF VANDERHAEGE'S TOAD-HEADED TURTLE, *Phrynops vanderhaegei* (Testudines, Chelidae) IN SE BRAZIL

FRANCO LEANDRO SOUZA^{1*}, MARCIO MARTINS² & RICARDO JANINI SAWAYA³

¹ Universidade Estadual de Campinas, Instituto de Biologia, Departamento de Parasitologia.
CP 6109. 13083-970, Campinas, SP. Brasil. e-mail: flsouza@unicamp.br

² Universidade de São Paulo, Instituto de Biociências, Departamento de Ecologia.
CP 11461. 05422-970, São Paulo, SP. Brasil. e-mail: jararaca@ib.usp.br

³ Universidade Estadual de Campinas, Instituto de Biologia, Departamento de Zoologia.
CP 6109. 13083-970, Campinas, SP. Brasil. e-mail: sawaya@unicamp.br

* corresponding author

Resumen: Se presenta una nueva localidad de la tortuga *Phrynops vanderhaegei* en Brasil sudoriental. Los especímenes se localizaron en un área de savana (*cerrado*) con pequeños tramos de bosque ripario. Se comenta brevemente el hábitat circundante. El hallazgo de esta población aislada puede ser importante en la conservación de la especie ya que indica que otras poblaciones podrían sobrevivir y reproducirse en pequeños fragmentos con vegetación bien conservada.

Key words: Brazil, Chelidae, conservation, *Phrynops vanderhaegei*, turtle.

Records related to geographical distribution for Brazilian freshwater turtles are relatively rare and the lack of such basic information has caused mistaken interpretations about species range (VANZOLINI, 1994). Since ecosystems are becoming highly fragmented, mainly due to human action, all information available should be used to identify areas that harbor threatened species, which would allow the development of appropriate management programs. In this note, we report a new locality record for the Vanderhaege's toad-headed turtle, *Phrynops vanderhaegei*, and provide comments on its conservation and habitat in southeastern Brazil.

Phrynops vanderhaegei is a medium-sized freshwater turtle, with carapace length around 250 mm (ERNST & BARBOUR, 1984; CABRERA, 1998). Few records concerning its natural history are available. Among these, anecdotal information indicates that this turtle is carnivorous and diurnal (ERNST & BARBOUR, 1989; CABRERA, 1998). Shallow water lagoons with dense aquatic vegetation and occasional scattered trees in the margins characterize its habitat (ERNST & BARBOUR, 1989; CABRERA, 1998). The distribution of *P. vanderhaegei* is still poorly known (Figure 1), with its southern limit at Mala Frigo (Santa Fé, Argentina) and the

northern limit at Cuiabá (Mato Grosso, Brazil; IVERSON, 1992; CABRERA, 1998).

For the state of São Paulo (Brazil), the only record for *P. vanderhaegei* is from a place around the cities of Perú and Caieiras




Figure 1. Map showing locality records of *Phrynops vanderhaegei* (modified from IVERSON, 1992). Star denotes the new record, Itirapina, São Paulo state, southeastern Brazil. Argentina: 1 - Mala Frigo; Paraguay: 2 - Asunción, 3 - Tobati, 4 - Nueva Germania, 5 - Capitán Brado, 6 - C. Curó; Brazil: 7 - Cuiabá, 8 - Perú/Caieiras, 9 - Itirapina.



Figure 2. A juvenile female of the Vanderhaege's toad-headed turtle, *Phrynops vanderhaegei*, captured at Itirapina, São Paulo state, southeastern Brazil Photo: R. J. Sawaya.

(IVERSON, 1992; IVERSON, *pers. comm.*), which is the easternmost known limit of its distribution (Figure 2). We found the Vanderhaege's toad-headed turtle at "Estação Ecológica de Itirapina" (Itirapina Ecological Station; hereafter Itirapina), which is located within the municipalities of Itirapina and Brotas (22°15' S, 47°49' W), São Paulo state, southeastern Brazil (Figure 1). This locality is about 180 km west of the Perú/Caieiras record. The Itirapina reserve is a 2300 ha tract of savannas (*cerrado*), from grasslands to dense arboreal formations, with small stretches of riparian forests (GIANOTTI, 1988; *pers. obs.*).

Three individuals of *P. vanderhaegei* (Figure 1) were captured in pitfall traps with drift fence (CECHIN & MARTINS, 2000), which are being used to survey the vertebrate fauna of Itirapina. Two were hatchlings (<1 month old; capture dates 5 and 25 February, 2000) and one was a juvenile female (capture date 3 March, 2000). Turtles were captured after some days of intense rainfall, which resulted in the flooding of the area. Their body measures are in Table 1.

Measures	hatchling 1	hatchling 2	juvenile female
SLC	39.2	41.9	148.7
CCL	44.7	44.4	160.2
CW	25.1	28.2	102.5
PL	28.7	33.1	128.3
PW	18.4	22.0	75.0
W	16.4	16.0	450.0

Table 1. Body measurements of three individuals of *Phrynops vanderhaegei* captured at Itirapina, São Paulo state, southeastern Brazil. Abbreviations are: straight line carapace length (SLC), curved line carapace length (CCL), carapace width (CW), plastron length (PL), plastron width (PW), and weight (W). All measurements are in millimeters; weight is in grams.

The habitat in which these turtles were captured is characterized by a small stretch of gallery forest (around 65 ha) surrounding a shallow, short tributary (about 1 km) of Lobo stream. Grasslands with a few shrubs (*campo sujo*) surround the gallery forest. The Lobo stream runs towards a large man made dam ("Represa do Lobo") about 1 km from the tributary where *P. vanderhaegei* was found. Besides the Itirapina reserve, few other natural habitats occur in the region; pastures and

commercial *Pinus* and *Eucalyptus* forests cover most of the area (GIANOTTI, 1988; pers. obs.).

In contrast with Itirapina and Cuiabá (Mato Grosso, Brazil), where the typical vegetation is composed of several savanna formations, the region of the Perús/Caieiras locality is covered by Atlantic rainforest (*sensu lato*). The Lobo stream belongs to the mid-Paraná river basin and the main river from Perús/Caieiras region, the Tietê river, is also a tributary of the Paraná river. The Tietê river runs through a wide area of the central region of São Paulo state, with its origin located in the west side of the Serra do Mar mountain complex (Figure 2). On the other hand, rivers where *P. vanderhaegei* were recorded in Cuiabá (Brazil), Paraguayan, and Argentinean localities belong to the Paraguay river basin. Hundreds of years ago, the vegetation along these river systems was continuous and both the coastal Atlantic rainforests and the upland mesophytic forests extended their domains to wide areas. However, due to human action beginning around the XVII century, these forests were drastically reduced (DEAN, 1996). Thus, species which once could make migrations throughout these vegetation corridors, became isolated in disjunct populations. The dependence of such corridors for migration is recorded for some groups, including reptiles, birds, and mammals (VANZOLINI, 1976; REDFORD & FONSECA, 1986; GUIX, 1997; VIVO, 1997). If the record of *P. vanderhaegei* for Perús/Caieiras region is correct, then probably the species could be once spread throughout São Paulo state, reaching also Mato Grosso. Turtles could have used rivers surrounded by riparian forests, even small ones like those of Itirapina.

Turtle species richness is high in Brazil (PRITCHARD & TREBBAU, 1984; IVERSON, 1992) and several species are often included in conservation programs (IUCN/SSC, 1991; IBGE, 1995; SEMA, 1998). Since species often exhibit specific habitat requirements, such as unpolluted streams and places with dense aquatic vegetation (PRITCHARD & TREBBAU, 1984; ERNST & BARBOUR, 1989; CABRERA, 1998), they are highly vulnerable to becoming restricted to isolated populations. This is an important concern because many turtles species inhabiting areas threatened by human action are

disappearing (ROCHA-E-SILVA & KISCHLAT, 1992). The *cerrado* vegetation covers about one fourth of Brazilian territory (ca. 2 million km²; Eiten, 1972, 1992; IBGE, 1993; RATTER *et al.*, 1997), but due to human action, mainly agricultural practices and charcoal production, the *cerrado* has experienced severe reductions in its area in the last decades. Presently, only 1.5% of this biome is preserved as Federal Reserves (IBGE, 1993; RATTER *et al.*, 1997). The finding of juvenile and hatchlings of *P. vanderhaegei* living in small savanna fragments like those of Itirapina, indicates that even small reserves can be important to the conservation of this species, with some populations surviving and reproducing in small patches.

Acknowledgments: The authors are grateful to F. B. Molina for early identification of the turtles. J. Iverson kindly permitted the use of the distribution map and provided locality records. The Instituto Florestal and D. Zancheta for logistical support; and L. Anjos, V. Bonato, and C.A. Brasileiro for field assistance. This is the paper number 2 of the project "Ecology of the *Cerrados* of Itirapina". Fieldwork was partially supported by a grant from FAPESP (95/09642-5) and additional funds from Universidade de São Paulo.

REFERENCES

- CABRERA, M. (1998): *Las Tortugas Continentales de Sudamérica Austral*. BR Copias, Córdoba, Rep. Argentina.
- CECHIN, S. Z. & MARTINS, M. (2000): Eficiência de armadilhas de queda (pitfall traps) em amostragens de anfíbios e répteis no Brasil. *Revta. brasil. Zool.*, 17: 729-740.
- DEAN, W. (1996): *A Ferro e Fogo: a História e a Devastação da Mata Atlântica Brasileira*. Cia. das Letras. São Paulo, SP.
- EITEN, G. (1972): The cerrado vegetation of Brazil. *Bot. Rev.*, 38: 201-341.
- EITEN, G. (1992): Natural brazilian vegetation types and their causes. *An. Acad. brasil. Ci.*, 64 (Supl. 1): 35-65.
- ERNST, C. H. & BARBOUR, R. W. (1989): *Turtles of the World*. Smithsonian Institution Press, Washington, D.C.
- GIANOTTI, E. (1988): *Composição florística e estrutura fitossociológica da vegetação de cerrado e mata ciliar da Estação Experimental de Itirapina (SP)*. Ms. Thesis. Universidade Estadual de Campinas. Campinas, SP.

- Guix, J. C. (1997): Exclusão geográfica e ecológica de *Penelope obscura*, *Penelope superciliaris* e *Pipile jacutinga* (Galliformes, Cracidae) no estado de São Paulo. *Ararajuba*, 5: 195-202.
- IBGE. (1993): *Mapa de Vegetação do Brasil*. Instituto Brasileiro de Geografia e Estatística, Rio de Janeiro, RJ.
- IBGE. (1995): *Mapa da Fauna Brasileira Ameaçada de Extinção*. Instituto Brasileiro de Geografia e Estatística, Rio de Janeiro, RJ.
- IUCN/SSC. (1991): *Tortoises and Freshwater Turtles. Na Action Plan for their Conservation*. Compiled by David Stubbs. Cambridge, U.K.
- IVERSON, J. B. (1992): *A Revised Checklist with Distribution Maps of the Turtles of the World*. Privately Printed, Richmond, Indiana.
- PRITCHARD, P. C. H. & TREBBAU, P. (1984): *The Turtles of Venezuela*. Society for the Study of Amphibians and Reptiles, Athens, Ohio.
- RATTER, J. A.; RIBEIRO, J. F. & BRIDGEWATER, S. (1997): The Brazilian cerrado vegetation and threats to its biodiversity. *An. bot. (London)*, 80: 223-230.
- REDFORD, K. H. & FONSECA, G. A. B. DA (1986): The role of gallery forests in the zoogeography of the cerrado's non-volant mammals fauna. *Biotropica*, 18: 126-135.
- ROCHA-E-SILVA, R. & KISCHLAT, E-E. (1992): Considerações sobre quelônios de água-doce no Estado do Rio de Janeiro. Pp. 1-17. in: *Second International Symposium on Environmental Studies of Tropical Rainforests*. Rio de Janeiro, RJ.
- SEMA. (1998): *Secretaria do Estado do Meio Ambiente. Fauna ameaçada no estado de São Paulo*. São Paulo, SMA/CED.
- VANZOLINI, P. E. (1976): On the lizards of a cerrado-caatinga contact: evolutionary and zoogeographical implications (Sauria). *Papéis Avulsos Zool., S. Paulo*, 29: 111-119.
- VANZOLINI, P. E. (1994): On the distribution of certain South American Turtles (Testudines: Testudinidae & Chelidae). *Smithsonian Herpetol. Inform. Service*, 97: 1-10.
- VIVO, M. DE (1997): Mammalian evidence of historical ecological change in the caatinga semiarid vegetation of northeastern Brazil. *J. Comp. Biol.*, 2: 65-73.
-