

Communal egg-laying and nest-sites of the Goo-eater Snake, *Sibynomorphus mikanii* (Dipsadidae, Dipsadinae) in southeastern Brazil

HENRIQUE B. P. BRAZ^{1, 3, 4}, FRANCISCO L. FRANCO²
and SELMA M. ALMEIDA-SANTOS¹

¹ Laboratório de Ecologia e Evolução, Instituto Butantan, 05503-900, São Paulo SP, Brasil

² Laboratório de Herpetologia, Instituto Butantan, 05503-900, São Paulo SP, Brasil

³ Programa de Pós-graduação Interunidades em Biotecnologia, Universidade de São Paulo, SP, Brasil

⁴ Corresponding author: hbraz@butantan.gov.br

FOR oviparous reptiles without parental behaviour, female nest-site selection plays a significant role in the evolution of life histories (Resetarits, 1996; Shine, 2004). Nevertheless, nest-sites and oviposition modes of neotropical snakes are relatively unknown, mostly because mothers are so successful at hiding their eggs that nests are rarely found in nature. Additionally, much emphasis has been placed on life history components such as body size, number and size of offspring, and age at maturity (Stearns, 1992; Resetarits, 1996).

Snakes oviposit under rocks, logs or any other surface cover, in preformed subterranean chambers (Packard & Packard, 1988) and within nests of other animals such as alligators (Hall & Meier, 1993), ants and termites (Riley et al., 1985). With regard to oviposition modes, snakes oviposit both in solitary and communal nests (Vaz-Ferreira et al., 1970; Graves & Duvall, 1995; Blouin-Demers et al., 2004).

Communal oviposition is a widespread phenomenon and occurs when several females, conspecifics or not, share the same nest cavity to deposit their eggs (Graves & Duvall, 1995). Among neotropical species, reports of nest-sites and communal nests are scarce and are mostly related to colder climates (e.g. Vaz-Ferreira et al., 1970; Cadle & Chuna, 1995). Recently, Albuquerque & Ferrarezzi (2004) reported one communal nest for the neotropical colubrid snake *Sibynomorphus mikanii* in an anthropized area in southeastern Brazil. Herein we describe another

three nest-sites, nesting areas and oviposition modes of the goo-eater snake *S. mikanii* in southeastern Brazil.

Sibynomorphus mikanii is a dipsadine snake that feeds on slugs (Laporta-Ferreira et al., 1986; Oliveira, 2001) and is distributed in Central, Atlantic and Meridional Brazilian uplands in cerrado and tropical forest areas (Franco, 1994). Females oviposit from September (early spring) to February (mid-summer) and clutch size varies from three to 10 eggs, averaging 5.9 (Oliveira, 2001).

On 5 February 2007, 41 eggs and 11 empty shells (Fig. 1c) were found together inside a hole, 20 cm below ground surface, at the edge of a degraded wood inside Instituto Butantan (IBSP), São Paulo city, Brazil. The nest (hereafter nest #1) was situated 17 m away from the wood and 9 m away from the backyard of a house on a slightly steep slope (Fig. 1a). A small hole (50 mm diameter), at the side of the nest may have provided access for the snakes (Fig. 1b). Temperature at the same depth around the nest averaged 27.3°C (range = 27-28°C). Four empty shells contained fluids indicating recent hatchings whereas seven were completely desiccated. Two dead hatchlings were found near the nest (± 1 m away). Nest #2 and #3 were discovered by a farmer in two different spots, in a 5000 m² house backyard, in Vargem city, Brazil. Nest #2 was discovered under dry grass accumulated after ground weeding and contained nine eggs. Nest #3 was found 30 m away from nest #2 under a large

rock (30 x 60 x 15 cm) and had a total of 12 eggs. Eggs were donated to IBSP on 1 March 2007, some days after collection and by this time one egg had hatched. In the laboratory, we counted four fresh empty shells and 17 eggs (three dehydrated and one parasitized by fungi). In both backyards, dogs circulated freely over the nesting areas. After the donation of the eggs, the farmer found a dead hatchling of *S. mikanii* near nest #3. Eggs were measured, weighed (Table 1), and incubated in a laboratory at 27°C (temperature of nest #1). Fungal infection precluded eight eggs (19.5%) from nest #1 and five eggs (45.5%)

nine different oviposition events may have occurred in nest #1, one to two in nest #2, and two in nest #3 in the current reproductive season. These assumptions are strengthened if we take into consideration hatching dates (seven in nest #1 and three in nest #2 and #3). Although communal oviposition in nest #1 and #3 is evident, there is some doubt over nest #2 as it is quite possible that one single female laid the nine eggs.

Nest-sites and nesting areas are described for few species of neotropical snakes (e.g. Vaz-Ferreira et al., 1970; Cadle & Chuna, 1995; Albuquerque & Ferrarezzi, 2004).




Figure 1. (A) Diagrammatic vertical section of the terrain surrounding nest #1; (B) view of the destroyed nest and its likely entrance; (C) Eggs of *Sibynomorphus mikanii* found within nest #1.

from nest #2 and #3 from hatching. Eggs were dissected but we were unable to find or to identify the embryos. One hatchling from nest #2 died half emerged from the eggshell. Thus, 33 successful hatchlings from nest #1 occurred between 6 February and 16 April 2007, whereas from nest #2 and #3 five occurred between 1 March and 31 May 2007. Hatchlings were measured, weighed, and sexed by eversion of hemipenis (Table 1). Based on mean clutch size of the species (5.9 eggs [Oliveira, 2001]), we inferred that nearly seven to

Despite the fact that some snakes dig a hole in the soil to oviposit (e.g. Burger & Zappalorti, 1986), most species apparently are unable to construct a nest and rely on pre-existing sites for oviposition (Packard & Packard, 1988). This seems to be the case for *S. mikanii* (Albuquerque & Ferrarezzi, 2004; this study). Although eggs of nest #1 were found inside a hole, it is unlikely that any female of *S. mikanii* actually excavated it because the soil was very compacted. Moreover, in the laboratory, gravid *S. mikanii* tend to hide the eggs under the

water bowl or under rocks instead of burying them (H.B.P. Braz & S.M. Almeida-Santos, unpublished data). In addition, other dipsadine snakes also oviposit in pre-existing sites (Brandão & Vanzolini, 1985; Riley et al., 1985; Cadle & Chuna, 1995; Greene, 1997).

Thermal conditions are often suggested as a factor driving maternal choice of nest-site (Blouin-Demers et al., 2004; Shine 2004) because incubation temperature affects offspring phenotypes (Deeming, 2004) and therefore may influence organismal fitness (Elphick & Shine, 1998; Brown & Shine, 2004). Females frequently oviposit in forest clearings (Fowler, 1966; Brodie et al., 1969; Covacevich & Limpus, 1972;

nests are usually higher than in solitary ones (e.g. Blouin-Demers et al., 2004) due to metabolic heat generated by embryos (Burger, 1976; Ewert & Nelson, 2003).

Communal nesting might be adaptive because higher temperatures in nests enhance hatchling phenotypes (Blouin-Demers et al., 2004). However, studies on the thermal and hydric requirements of *S. mikanii* embryos would be needed to test these hypotheses. In parallel, egg aggregations also offer other potential advantages such as protection (Graves & Duvall, 1995; Jackson, 1998) and predator satiation (Eckrich & Owens, 1995; Graves & Duvall, 1995). If communal oviposition offers such advantages to hatchlings (e.g. phenotype

Measurements	Nest #1	Nest #2 and #3
Eggs	n = 41	n = 13
Length (mm)	27.9 ± 2.7	25.7 ± 3.0
Width (mm)	14.5 ± 2.0	12.6 ± 0.9
Mass (g)	3.5 ± 0.9	2.5 ± 0.4
Hatchlings	n = 33	n = 5
SVL (mm)	171.8 ± 13.5	170.0 ± 5.8
TL (mm)	34.0 ± 4.7	29.4 ± 2.6
Mass (g)	2.3 ± 0.4	2.0 ± 0.5
Sex (male/female)	19/14	2/3

Table 1. Measurements of eggs and hatchlings of three natural nests of the Goo-eater Snake *Sibynomorphus mikanii*. SVL = Snout-vent length; TL = Tail length.

Burger & Zappalorti, 1986; Albuquerque & Ferrarezzi, 2004) and nests located in these areas generally are hotter than nests located in shaded areas because shading reduces insolation and heating of the soil (Magnusson & Lima, 1984; Shine et al., 2002). As well as nest #1, several nests have also been found in slopes (e.g. Brodie et al., 1969; Covacevich & Limpus, 1972; Burger, 1976; Albuquerque & Ferrarezzi, 2004; James & Henderson 2004) and factors like direction and slope influence the absorption of solar radiation (Burger, 1976). Thus, mothers may have selected these sites in seeking to maximize sunlight exposure to accelerate embryonic development or optimize phenotypic traits of the resulting hatchlings. Therefore thermal conditions could also be a major factor influencing communal nesting behaviour. Temperatures in communal

improvement, predator satiation), why, then, would one female oviposit in a solitary nest as is likely to have occurred in nest #2? Blouin-Demers et al. (2004) suggested that the disadvantages of solitary nests may be compensated by lower risk of egg parasitism by fungi. Additionally, in communal nests availability of water is less than in solitary nests (Marco et al., 2004; Radder & Shine, 2007). This modifies hydric exchange between the eggs and the environment and the consequences to hatchling phenotypes may be more detrimental to aggregated eggs (Marco et al., 2004). Thus it is reasonable to suggest that there are trade-offs between these two modes of egg-laying that result in similar fitness payoffs (Blouin-Demers et al., 2004).

In summary, there are two (but nonexclusive) reasons for the occurrence of communal nesting

behaviour: scarcity of suitable nesting sites (e.g. optimum moisture and temperature; protection against predators) or adaptive behaviour; that increases reproductive success due to aggregation in large clusters.

Our findings plus literature data indicate a preference of gravid *S. mikanii* to nest communally even when similar potential nest-sites were present in nesting areas. We suggest that such widespread behaviour might result from adaptation. However, the adaptive significance of communal oviposition remains unknown.

ACKNOWLEDGEMENTS

We thank D. A. C. Martins, and the staff of Vigilância Sanitária de Vargem for the material donation and information. J. C. Ferreira and A. C. Barbosa for laboratory assistance. B. M. Graves and D. W. Owens kindly provided references. This study was supported by FAPESP (07/51977-3).

REFERENCES

- Albuquerque, C.E. & Ferrarezzi, H. (2004). A case of communal nesting in the Neotropical snake *Sibynomorphus mikanii* (Serpentes, Colubridae). *Phyllomedusa* **3**, 73-77.
- Blouin-Demers, G., Weatherhead, P.J. & Row, J.R. (2004). Phenotypic consequences of nest-site selection in Black Rat snakes (*Elaphe obsoleta*). *Can. J. Zool.* **82**, 449-456.
- Brandão, C.R.F. & Vanzolini, P.E. (1985). Notes on incubatory inquilinism between Squamata (Reptilia) and the Neotropical fungus-growing ant genus *Acromyrmex* (Hymenoptera: Formicidae). *Papeis Avulsos Zool.* **36**, 31-36.
- Brodie, E.D., Jr., Nussbaum, R.A. & Storm, R. M. (1969). An egg-laying aggregation of five species of Oregon Reptiles. *Herpetologica* **25**, 223-227.
- Brown, G.P. & Shine, R. (2004). Maternal nest-site choice and offspring fitness in a tropical snake (*Tropidonophis mairii*, Colubridae). *Ecology* **85**, 1627-1634.
- Burger, J. (1976). Temperature relationships in nests of the northern diamondback terrapin, *Malaclemys terrapin terrapin*. *Herpetologica* **32**, 412-418.
- Burger, J. & Zappalorti, R.T. (1986). Nest site selection by pine snakes, *Pituophis melanoleucus*, in the New Jersey Pine Barrens. *Copeia* **1986**, 116-121.
- Cadle, J.E. & Chuna, P.M. (1995). A new lizard of the genus *Macropholidus* (Teiidae) from a relictual humid forest of northwestern Peru, and notes on *Macropholidus ruthveni* Noble. *Breviora* **501**, 1-39.
- Covacevich, J. & Limpus, C. (1972). Observations on community egg-laying by the yellow-faced whip snake, *Demansia psammophis* (Schlegel) 1837 (Squamata: Elapidae). *Herpetologica* **28**, 208-210.
- Deeming, D.C. (2004). Post-hatching phenotypic effects of incubation in reptiles. In: *Reptilian Incubation: Environment, Evolution, and Behaviour*. pp. 229-251, D.C. Deeming (Ed.). Nottingham: Nottingham University Press.
- Eckrich, C.E. & Owens, D.W. (1995). Solitary versus arribada nesting in the olive ridley sea turtles (*Lepidochelys olivacea*): a test of the predator-satiation hypothesis. *Herpetologica* **51**, 349-359.
- Elphick, M.J. & Shine, R. (1998). Longterm effects of incubation temperature on the morphology and locomotor performance of hatchling lizards (*Bassiana duperreyi*, Scincidae). *Biol. J. Linn. Soc.* **63**, 429-447.
- Ewert, M.A. & Nelson, C.E. (2003). Metabolic heating of embryos and sex determination in the American alligator, *Alligator mississippiensis*. *J. Therm. Biol.* **28**, 159-165.
- Fowler, J.A. (1966). A communal nesting site for the smooth green snake in Michigan. *Herpetologica* **22**, 231.
- Franco, F.L. (1994). O gênero *Sibynomorphus* Fitzinger, 1843 no Brasil (Colubridae: Xenodontinae, Dipsadinae). Unpublished M. Sc. Dissertation. Pontifícia Universidade Católica do Rio Grande do Sul, Brazil.
- Graves, B.M. & Duvall, D. (1995). Aggregation of squamate reptiles associated with gestation, oviposition, and parturition. *Herpetol. Monogr.* **9**, 102-129.
- Greene, H.W. (1997). *Snakes: the Evolution of Mystery in Nature*. Berkeley: University of California Press.

- Hall, P.M. & Meier, A.J. (1993). Reproduction and behaviour of western mud snakes (*Farancia abacura reinwardtii*) in american alligator nests. *Copeia* **1993**, 219-222.
- Jackson, D. (1989). Turtles use of alligator nests. In: *Crocodyles and Alligators*. pp. 145. C. A. Ross (Ed.). London: Mercurst Press.
- James, A. & Henderson, R.W. (2004). Communal nesting site in the snake *Liophis juliae* in Dominica, West Indies. *Carib. J. Sci.* **40**, 263-265.
- Laporta-Ferreira, I.L., Salomão, M.G. & Sawaya, P. (1986). Biologia de *Sibynomorphus* (Colubridae-Dipsadinae) – Reprodução e hábitos alimentares. *Rev. Bras. Biol.* **46**, 793-799.
- Magnusson, W.E. & Lima, A.P. (1984). Perennial communal nesting by *Kentropyx calcaratus*. *J. Herpetol.* **18**, 73-75.
- Marco, A., Díaz-Paniagua, C. & Hidalgo-Vila, J. (2004). Influence of egg aggregation and soil moisture on incubation of flexible-shelled lacertid lizard eggs. *Can. J. Zool.* **82**, 60-65.
- Oliveira, J.L. (2001). Ecologia de três espécies de dormideira, *Sibynomorphus* (Serpentes: Colubridae). Unpublished M.Sc. Dissertation. Universidade de São Paulo, Brazil.
- Packard, G.C. & Packard, M.J. (1988). The physiological ecology of reptilian eggs and embryos. In: *Biology of the Reptilia, Vol. 16. Ecology B, Defense and Life History*, pp. 523-606. C. Gans & R. B. Huey (Eds.). New York: C. Alan R. Liss Inc.
- Radder, R.S. & Shine, R. (2007). Why do female lizards lay their eggs in communal nests? *J. Anim. Ecol.* **76**, 881-887.
- Resetarits, W.J., Jr. (1996). Oviposition site choice and life history evolution. *Amer. Zool.* **36**, 205-215.
- Riley, J., Stimson, A.F. & Winch, J.M. (1985). A review of Squamata ovipositing in ant and termite nests. *Herp. Rev.* **16**, 38-43.
- Shine, R. (2004). Adaptive consequences of developmental plasticity. In: *Reptilian Incubation: Environment, Evolution, and Behaviour*, pp. 187-210. D. C. Deeming (Ed.). Nottingham: Nottingham University Press.
- Shine, R., Barrott, E.G. & Elphick, M.J. (2002). Some like it hot: effects of forest clearing on nest temperatures of montane reptiles. *Ecology* **83**, 2808-2815.
- Stearns, S.C. (1992). *The Evolution of Life Histories*. Oxford: Oxford University Press. 249 pp.
- Vaz-Ferreira, R., Zolessi, L.C. & Achával, F. (1970): Oviposicion y desarrollo de Ofidios y Lacertilios em hormigueiros de *Acromyrmex*. *Physis* **29**, 431-459.