

ACKNOWLEDGMENTS

We thank M. Harvey, K. Kittelson, and T. Dorcas for assistance in the field. J. A. Campbell provided helpful comments on the manuscript and logistical assistance. J. L. Darling curated the specimens expeditiously. This work was authorized by the Texas Parks and Wildlife Dept. (permit number: SPR-0590-072).

LITERATURE CITED

- Dixon, J. R. 1987. Amphibians and Reptiles of Texas. Texas A&M University Press, College Station. 434 pp.
- Mecham, J. S. 1983. *Nerodia harteri*. Cat. Am. Amph. Rept. 330.1-330.2.
- Scott, N. J., Jr., T. C. Maxwell, O. W. Thornton, Jr., L. A. Fitzgerald, and J. W. Flury. 1989. Distribution, habitat, and future of Harter's water snake, *Nerodia harteri*, in Texas. J. Herpetol. 23(4):373-389.
- Smith, D. D. 1983. Geographic distribution: *Nerodia harteri harteri*. Herpetol. Rev. 14:84-85.
- Trapido, H. 1941. A new species of *Natrix* from Texas. Amer. Midl. Natur. 32:673-680.
- Wade, V. E. 1968. A range extension of the water snake, *Natrix harteri harteri* Trapido. Texas J. Sci. 20:194-196.

MICHAEL E. DORCAS¹
JOSEPH R. MENDELSON III²
 Department of Biology
 UTA Box 19498
 University of Texas at Arlington
 Arlington, Texas 76019, USA.

¹Current Address:
 Department of Biological Sciences
 Campus Box 8007
 Idaho State University
 Pocatello, Idaho 83209-0009, USA

²Current Address:
 Museum of Natural History
 University of Kansas
 Lawrence, Kansas 66045-2454, USA.

PREDATION ON AVIAN EGGS BY THE BOID SNAKE, *Eunectes notaeus*

Besides the well-known, specialized egg-eater *Dasypeltis* (Gans 1974), several colubrid snakes (including *Boiga*, *Conopsis*, *Elachistodon*, *Elaphe*, *Lampropeltis*, *Pseustes*, and *Spilotes*) include bird eggs in their diets (Cunha and Nascimento 1978, Greene 1989, Mushinsky 1987, Savitzky 1983, Scott 1983). To date, documented observations on avian egg-eating seem restricted to colubrid snakes.

Constricting anacondas, boid genus *Eunectes*, are known to mainly feed on avian and mammalian prey (Cunha and Nascimento 1978, Ditmars 1912, Wehekind 1955), and occasionally on crocodylians and turtles (Parker and Grandison 1977, Yamashita et al. 1985). In southwestern Brasil we observed the yellow anaconda, *E. notaeus*, feeding on prey as diverse as fishes, turtles, caimans, wading birds, and rodents; this snake even scavenges, a habit anticipated by Sazima and Strüssmann (1990). Here we report on an observation of avian egg-eating by *E. notaeus*.

Field work was done in the Pantanal wetlands at Poconé, Mato Grosso, southwestern Brasil. The Pantanal covers ca. 100,000 km² of almost flat, seasonally flooded terrain (additional information in Prance and Schaller 1982). As part of a study on the Pantanal herpetofauna, 33 anacondas were caught and palpated to determine whether their stomach contained food. Whenever a bolus was detected, the snake was forced to regurgitate its prey (Slip and Shine 1988), which was then examined.

A juvenile *E. notaeus* (165 cm SVL, 3.2 kg) was found on 11 July 1990, apparently basking on an emergent mat of aquatic vegetation at 1745 h, with an air temperature of 19.5°C. Upon manipulation the snake regurgitated two intact bird eggs; an additional egg was broken during palpation. The eggs were identified as those of limpkin (Harrison 1978), *Aramus guarana*, a common wading bird in the Pantanal.

Head length and head width of the snake were 5.4 cm and 3.3 cm, respectively. The two intact eggs measured 6.2 x 4.3 cm and ca. 6.0 x 4.5 cm, and weighed about 60 g each, a prey/predator mass ratio (Pough and Groves 1983) of ca. 0.02. This is a very low mass food item for a boid snake (average of 0.36; see Greene 1983).

The eggs were all swallowed by their pointed end first, an orientation commonly found in other instances of egg-eating by snakes (see Gans 1974). Egg handling by *E. notaeus* may possibly be similar to that employed by the boid *Loxocemus bicolor* when feeding on reptile eggs—either pushing the egg against the body using its mouth, or enveloping the eggs with body loops (Mora 1987). A young yellow anaconda was observed to engulf small fishes using the first technique (A. S. Abe, pers. comm.).




Figure 1. The 100 km hiatus in the distribution of *Nerodia h. harteri* along the Brazos River reported by Scott et al. (1989). Closed circles represent localities where snakes were found by Scott et al. (1989). Lettered localities are referred to in text.

Feeding on avian eggs is regarded as a kind of durophagy (Savitzky 1983), and anacondas also take turtles (Parker and Grandison 1977), another hard-shelled prey ingested intact. The snake-necked turtle, *Platemys macrocephala*, was reported from intestinal contents of *E. notaeus* in the Pantanal (Yamashita et al. 1985), and we observed the turtle *Phrynops vanderhaegei* in the contents of an *Eunectes* rotting carcass (probably *E. murinus*) at another locality in Mato Grosso.

Limpkin nests may be found on bushes up to 1 m above water level in flooded terrain, or near the water in low trees up to 2 m (pers. obs.). Anacondas are known to occasionally climb (Parker and Grandison 1977), and we observed *E. notaeus* foraging at night and taking birds perched on bushes and trees at heights of about 2.5 m. Large nesting colonies of herons and ibises are a common sight in the Pantanal (see Yamashita and Valle 1990), and at these

places arboreality would be advantageous for the anaconda. Nestlings and eggs found by *E. notaeus* foraging in trees probably fall prey to this apparently generalized predator.

ACKNOWLEDGMENTS

We thank A. S. Abe, C. Gans, and M. Martins for critically reading an earlier draft of the manuscript; O. Cid at the Santa Inês Ranch for help in the fieldwork and logistical support; the CNPq provided financial support for fieldwork (grants 300092/79 to IS and 831105/89-5 to CS).

LITERATURE CITED

- Cunha, O. R., and F. P. Nascimento. 1978. Ofídios da Amazônia. X- As cobras da região leste do Pará. Mus. Para. Emílio Goeldi Publ. Avulsas no. 31.
- Ditmars, R. L. 1912. The feeding habits

AN INVITATION TO JOIN THE HERPETOLOGISTS' LEAGUE

The Herpetologists' League, established in 1936, is an international organization of people devoted to studying the biology of amphibians and reptiles. HL publishes two excellent scholarly journals: the quarterly *Herpetologica*, containing original research papers, essays and book reviews; and the annual supplement *Herpetological Monographs*, containing longer research articles, syntheses, and reviews that can be accommodated in a quarterly journal. The 1992 issue of HM will be a special expanded edition to accommodate publication of a recent symposium entitled "Amphibian Relationships: Phylogenetic Analysis of Morphology and Molecules." Both journals are mailed free of charge to members in good standing of the Herpetologists' League. HL currently provides airlift delivery of its journals (except back issues) to countries outside the United States at no extra charge.

The Herpetologists' League recently embarked on a program to support the publication of books of interest to its members, and to provide members the opportunity to purchase these books for a limited time at specially discounted prices. The initial result of this program was the critically acclaimed "Venomous Reptiles of Latin America." Future projects in this series include the herpetofauna of Baja California, as well as continuing the world checklists project started by the Association of Systematics Collections. Volumes currently in progress include one on snakes and a revision of the 1985 amphibian checklist.

To receive an application for membership, write to:


Dr. Andrew H. Price
Texas Natural Heritage Program
Texas Parks and Wildlife Department
4200 Smith School Road
Austin, Texas 78744, USA

CLOACAL QUICK-READING THERMOMETERS

(Formerly Schultheis)

0/50°C

CALL OR WRITE TO
PRESENT MANUFACTURER



MILLER & WEBER
Inc.

Dept. H

1637 George Street
Ridgewood, Queens
New York, 11385

718-821-7110

FAX: 718-821-1673

Mfg. Since 1941

of serpents. *Zoologica* (NY) 1:197-238.

Gans, C. 1974. *Biomechanics: An Approach to Vertebrate Biology*. J. P. Lippincott, Philadelphia. 261 pp.

Greene, H. W. 1983. Dietary correlates of the origin and radiation of snakes. *Amer. Zool.* 23:431-441.

_____. 1989. Ecological, evolutionary, and conservation implications of feeding biology in Old World cat snakes, genus *Boiga* (Colubridae). *Proc. California Acad. Sci.* 46:193-207.

Harrison, C. 1978. *A Field Guide to the Nests, Eggs and Nestlings of North American Birds*. Collins, Glasgow.

Mora B., J. M. 1987. Predation by *Loxocemus bicolor* on the eggs of *Ctenosaura similis* and *Iguana iguana*. *J. Herpetol.* 21:334-335.

Mushinsky, H. R. 1987. Foraging ecology. In: R. A. Seigel, J. T. Collins, and S. S. Novak (eds.), *Snakes: Ecology and Evolutionary Biology*, pp. 302-334. MacMillan Publ. Co., New York.

Parker, H. W., and A. G. C. Grandison. 1977. *Snakes: A Natural History*. British Museum (Nat. Hist.) and Cornell Univ. Press, London and Ithaca. 108 pp.

Pough, F. H., and J. D. Groves. 1983. Specializations of the body form and food habits of snakes. *Amer. Zool.* 23:443-454.

Prance, G. T., and G. B. Schaller. 1982. Preliminary study of some vegetation types of the Pantanal, Mato Grosso, Brazil. *Brittonia* 34:228-251.

Savitzky, A. H. 1983. Coadapted character complexes among snakes: fossoriality, piscivory, and durophagy. *Amer. Zool.* 23:397-409.

Sazima, I., and C. Strüssmann. 1990. Necrofagia em serpentes brasileiras: exemplos e previsões. *Rev. Brasil. Biol.* 50:463-468.

Scott, N. J. 1983. *Conophis lineatus* (guardacamino). In: D.H. Janzen (ed.), *Costa Rican Natural History*, pp. 392-393. Univ. Chicago Press, Chicago.

Slip, D. J., and R. Shine. 1988. Feeding habits of the diamond python, *Morelia s. spilota*: ambush predation by a boid snake. *J. Herpetol.* 22:323-330.

Wehckind, L. 1955. Notes on the foods of the Trinidad snakes. *Brit. J. Herpetol.* 2:9-13.

Yamashita, C., J. C. Dalponte, M. P. Vale, and M. Friedlander. 1985. Miscelânea sobre Alligatoridae e Chelidae (Reptilia) em Mato Grosso. *Resumos XII Congr. Bras. Zool. Campinas*:238-239 (abstract).

_____, and M. P. Valle. 1990. Sobre ninhas de aves do Pantanal do município de Poconé, Mato Grosso, Brasil. *Vida Silv. Neotrop.* 2:59-63.

CHRISTINE STRÜSSMANN¹

Fundação Estadual do Meio Ambiente
78000 Cuiabá, Mato Grosso, Brasil
and

IVAN SAZIMA

Departamento de Zoologia
Universidade Estadual de Campinas
13081 Campinas, São Paulo, Brasil

¹Current address: Caixa Postal 1430
78000 Cuiabá, Mato Grosso, Brasil

**THE IWAKUNI
SHIROHEBIS,
A GROUP OF ALBINO
*Elaphe climacophora***

Albinism in wild populations of snakes is rare or uncommon (Bechtel and Bechtel 1981; Dyrkacz 1981; Hensley 1959; see Satoh and Miyamoto (1987) for records of albino snakes in Japan). The Iwakuni shirohebis are a group of

albino *Elaphe climacophora* occurring in a limited area of the city of Iwakuni (34°10'N, 132°13'E), Japan. As far as we know, this is the only instance among snakes where albinism has become established in a wild population. Due to the rarity of this phenomenon, the habitat known to support albino snakes was designated as a natural monument by the Japanese government in 1924. In 1972, the subject of designation was changed to the snakes themselves. Although the Iwakuni shirohebi has been briefly mentioned in English-language literature (Goris and Naganuma 1979; Maki 1931), the accounts were brief and little information was provided concerning the status of the snakes. This paper summarizes the past and present status of this group of snakes.

Among several types of color abnormalities described (Bechtel 1978; Bechtel and Bechtel 1985), the Iwakuni shirohebi consists of amelanistic albinos. All adult snakes are white to yellow. They have golden irises, red pupils, and red tongues. Juveniles have orange blotches on the dorsal and lateral surfaces of the body. The pattern of the blotches is the same as that of wild type juveniles. As the crossing of the albino snakes always produces albino snakes, the gene for albinism is presumed to be allelic.

Little information on historical abundance and distribution exists. In 1924, the residents of Marifu village, one of the predecessors of Iwakuni city, were questioned as to the number of snakes in each block of the village (Yamaguchi Prefecture 1925). The number of snakes was estimated as "no fewer than 1000" and the estimated distribution was about 400 ha (Yamaguchi Prefecture 1925; Watasé 1926). Habitats occupied by snakes were reported to be houses, storehouses, river banks, and stone walls. Active snakes were observed in the streets, gardens, and fields (Watasé 1926). It was not rare to see the albino snakes in human residences (Yamaguchi Prefecture 1925).

Since the 1970s, the city of Iwakuni has appealed to its citizens to report occurrences of albino snakes. The number of snakes reported is declining; the average number per year (followed by SD and range) was 22.6 (12.3, 12-40), 8.8 (3.6, 4-12) and 4.8 (3.0, 0-8) for 1974-1978, 1981-1985, and 1986-1990, respectively.

The authorities and citizens of Iwakuni city are concerned about the decreasing

**HERPETOLOGICAL BOOKS
& PUBLICATIONS FOR SALE**

SEND FOR MY 1990-91 CATALOG OF OVER 3800 HERP TITLES (\$2 for postage) AND MY NEW 1991-92 ADDITIONAL HERP TITLES (sent free upon request).

SPECIALS OF THE MONTH:

BATRACHIA OF NORTH AMERICA BY E.D. COPE 522 pp. 1966 REPRINT \$16.00 ppd.

SET OF FIVE POSTERS, FIRST PUBLISHED BY THE ARMY FOR EASY IDENTIFICATION & HABITS OF VENOMOUS SNAKES OF THE WORLD. \$9.00 ppd.

SEND TO:

HERPETOLOGICAL SEARCH SERVICE AND EXCHANGE
117 E. SANTA BARBARA RD.
LINDENHURST, N.Y. 11757

number of albino snakes. The city of Iwakuni and the Society for the Conservation of the Iwakuni shirohebi, organized by volunteers, are making great efforts to conserve the shirohebi. To breed the snakes, they constructed an indoor breeding facility (total floor space of 240 m²) and three outdoor pens (total space of 1560 m²). Financial aid from the Japanese government was obtained for construction of one of the outdoor enclosures. In general, the adult snakes are allowed to breed at random in the pens, and emerging hatchlings are collected and raised in the indoor facility. Juveniles are released into the outdoor pens after three years. Approximately 400 snakes are maintained in these facilities.

The normal type of *E. climacophora* is abundant in the city. Studies are under way in an attempt to elucidate the circumstances surrounding the establishment and maintenance of high-frequency albinism in *E. climacophora*.

This study was sponsored by Iwakuni city and the Society for the Conservation of the Iwakuni shirohebi.