

Predation by the Opossum *Didelphis marsupialis* on the Rattlesnake *Crotalus durissus*

SELMA MARIA ALMEIDA-SANTOS¹, MARTA MARIA ANTONIAZZI²,
OSVALDO AUGUSTO SANT'ANNA³, AND CARLOS JARED^{2*}

¹ Laboratory of Herpetology, Instituto Butantan, Av. Vital Brazil 1500, 05503-900, São Paulo, Brazil

² Laboratory of Cell Biology, Instituto Butantan, Av. Vital Brazil 1500, 05503-900, São Paulo, Brazil

³ Laboratory of Immunogenetics, Instituto Butantan, Av. Vital Brazil 1500, 05503-900, São Paulo, Brazil

Abstract: Opossums are considered natural predators of snakes and possess resistance to the venom of some viperids. The resistance of *Didelphis* to *Crotalus* venom has been demonstrated through biochemical and immunological assays. However, systematic observations on the behavior of adult *Didelphis* preying on venomous snakes have never been conducted. In this study the predatory and defensive behaviors of *Didelphis marsupialis* and *Crotalus durissus*, respectively, were analyzed in captivity. Defensive strategies showed by snakes included immobility, flight attempts, coiling, cocking, rattling, and counterattack with strikes and bites. The most common defensive behavior of the rattlesnakes was immobility. The way the opossums attacked was classified in three categories, depending on the defensive reactions presented by the snakes. On all occasions when the opossums were bitten, the injection of venom apparently did not affect the predation. The great ability in capturing and handling *Crotalus durissus* together with the apparent great tolerance to the venom shown by *Didelphis marsupialis* when preying on these snakes confirms the existent biochemical and immunological data about the resistance of opossums to crotalic venoms. In this way our data strongly reinforce the supposition that this species is an effective snake predator in nature.

Key words: Marsupialia; Behavior; *Didelphis*; Venom; *Crotalus*

INTRODUCTION

Didelphis marsupialis is a very common marsupial in Brazil, living mainly in forests

(Cerqueira, 1985). It is found in environments modified by humans, adapts well in urban areas, and is nocturnal, scansorial, and omnivorous. Its diet is composed of fruits, nectar, small vertebrates, and invertebrates (Emmons, 1990). Many reports cite opossums as natural predators of snakes including venomous species (Silva Jr., 1956; Fitch, 1960; Cordero and Nicolas,

* Corresponding author. Tel/Fax: +55 11 257-8905.

E-mail address: carlosjared@uol.com.br (C. Jared)

1986; Sazima, 1992). In addition, Brodie III (1993) observed attacks of opossums towards snake replicas in the field. At the park of Butantan Institute, São Paulo, Brazil, attacks by native opossums on the outdoor enclosure snakes have been often reported (W. Fernandes, pers. commun.).

A number of studies is available demonstrating that opossums are immune to the venom of some viperids (Domont et al., 1991). Besides the well-known immunity against *Bothrops* venom (Moussatché et al., 1979, 1990; Perales et al., 1994; Neves-Ferreira et al., 1997), resistance of *Didelphis* to the venoms of *Crotalus durissus* (Vellard, 1945; Moussatché et al., 1979, 1990) and *C. atrox* (Werner and Vick, 1977; Perez et al., 1979) has been observed. All these data were obtained from biochemical and immunological assays. Studies about the zoological implications of this resistance have never been undertaken.

Information about the predatory behavior of American marsupials is scarce. Sazima (1992) made a single observation on the behavior of *Didelphis* toward its prey; his work, however, was not specific to predation. Jared et al. (1998) made a few preliminary observations on young opossums attacking and killing young *Bothrops jararaca* in captivity. Despite the strong supposition about opossums being effective predators of viperids, experimental results demonstrating this fact do not yet exist.

On the other hand, although snakes present the most elaborate antipredator mechanisms hitherto described among reptiles, few papers are found about the defensive behavior of these animals toward their effective predators (Greene, 1973, 1988). Among venomous snakes, rattlesnakes are a differentiated group possessing a unique structure, the rattle, specialized in sonorous defensive signaling (Greene, 1988). According to Duvall et al. (1985) and Greene (1988), in nature the whole set of defensive behaviors in rattlesnakes corresponds to an increase of aggressiveness comprising

procrypsis (immobility associated with a cryptic coloration pattern), flight, body coiling, cocking (retracting of the coiled body and intimidation with strikes), rattling, head hiding, strikes and bites, and finally emptying of the anal glands. This set of behaviors or, in most cases, the combination of some of them, is sufficient to prevent the snakes from being killed by predators.

This paper describes a behavioral experiment in the laboratory where *Crotalus durissus* was offered to *Didelphis marsupialis*. It aims to understand the defensive and attack strategies of both animals. It also has the intention of comparing the obtained behavioral data with the existent biochemical and immunological information about the resistance of *Didelphis* to *Crotalus* venom. Drawing on these results, this work finally tries to make a few inferences about the predatory and defensive behaviors of these animals in nature.

MATERIALS AND METHODS

Animals

Healthy adults *D. marsupialis* (N = 12) collected from the woods of Butantan Institute and from the surroundings of São Paulo city were used. They weighed 1.5 to 2 kg.

The opossums were maintained in individual tanks measuring 0.80 × 0.70 m and 0.70 m in depth, and closed with a wire netting lid. A wooden box measuring 0.27 m in width, 0.34 m in length, and 0.20 m in height with an inclined wire netting front door was placed inside the tank for shelter. Due to its design, the box did not allow the opossum in the shelter to see the environment inside the tank. This shelter box occupied 16% of the total area of the tank and was removed daily from the tanks for cleaning, with the animal inside. After that, the tanks were lined with cardboard, and the box was replaced and reopened. Water and food (fruits and

minced meat) were supplied daily. Opening and closing of the box door were done with a hook-tipped stick.

Farmers from the State of São Paulo (Brazil) supplied the snakes to the Herpetology Laboratory of Instituto Butantan. Thirty-six healthy adult *Crotalus durissus* (total length = 60 to 90 cm) were used. They were maintained in captivity for at least 20 days without being fed before the experiments in order to guarantee that the venom glands were full of secretion.

Behavioral evaluation

The observations were conducted during a two-year period. Before initiating the experiments each animal was kept in captivity at least for a month for acclimation to the new environment. Each opossum was tested with one snake every 15 days for three times per opossum. Before each observation the opossums were fasted for two days. The experiments were conducted in darkness at night, the period when the animals are active. The tank was lighted inside with a 15 W lamp and closed with a glass plate instead of the netting lid. With this system the opossums were kept acoustically and visually isolated from the outside environment. At the same time, the inclined door of the box made it possible for the observers to see the opossum inside the box from top, through the glass plate. The animal was kept in this condition for 1 hr before the snake was offered. The system allowed the option for the opossum of attacking the snake or remaining inside the box. Observations were started as soon as a snake was gently placed into the tank. Twenty-five out of a total of 36 experiments were recorded on video tape and photographed. In the experiments where predation did not occur the total time of observation was 120 minutes. Five control experiments were conducted placing a snake alone into the tank and recording its behavior on video tape for 1 hr.

A binomial test was used to statistically

test the preference for tail attack. The null hypothesis was $p = 0.5$.

RESULTS

The opossum, even at night, constantly remained inside the shelter box. When food was placed in the tank, the opossum immediately started to smell it with visible movements of its snout and, in most cases, came out of the box to feed on it. Sometimes, however, after smelling the food for a few minutes it remained inside the box.

Two distinct phases of behavioral interactions between the opossum and the rattlesnake were characterized after the snake was placed in the tank: 1) behaviors before the attack, including the observations before the opossum left the box and 2) the attack itself, that was defined here as the set of behaviors observed from the moment the opossum came out of the box and approached the snake until effective predation occurred.

Before the attack, the opossum, inside the box, immediately smelled the snake for a few seconds to 2 min. During this time different reactions of the snake could be observed: 1) calm movements around the tank, 2) a quiet coiled posture, 3) immobility (freezing), 4) flight, 5) threatening coiled posture (cocking) or 6) rattling. The opossum, after smelling the snake, sometimes remained in the box, not leaving to catch it. In four experiments, when the opossum inside the box spent more time smelling the snake, rattling was observed; in all these occasions the opossum chose to remain in the box, not attacking the snake.

When the opossum attacked, it came out of the box, approached the snake quickly (Fig. 1A), and captured it (Fig. 1B). On three occasions, the attack was so quick that the snake had no time to react. Most times, however, the snake showed some type of reaction.

Three types of behavioral interactions were recognized during the attacks. Figure




FIG. 1. One of the possible sequences of predatory behaviors of the opossum on the rattlesnake. A: The opossum directly approaches the snake, which remains immobile. B: The opossum captures the snake by the tail. The snake remains passive, flicking the tongue. C: The opossum continues eating the snake from the tail. The snake still flicks its tongue. D: The snake suddenly bites the opossum. E: The opossum reacts, kills the snake by chewing its head (arrow) and bites the rest of the body. F to H: The opossum continues eating the snake from the anterior end. While eating, it remains in the same position (F, G) until it finishes the whole snake (H).

2 summarizes the feeding tactics of the opossums, which depended on the reactions exhibited by the snakes after the initial approach:

- (1) The snakes showed an erratic behavior, moving the body by chance in a disoriented manner. In this case, the opossum immobilized them with successive bites along the whole body. Then it consumed them from either the tail or the head.
- (2) The snakes were immobile. In this case the opossum consumed them alive, preferentially from the tail (Fig. 1C). Usually the snakes stayed immo-

bile throughout predation. Sometimes, however, the snakes counterattacked with part of their bodies already eaten.

- (3) The snakes counterattacked with strikes and bites either just before being captured or having already part of its body eaten from the tail (Fig. 1D). In both cases the opossum killed them by chewing the head, and then immediately consumed them beginning from one of the extremities (Fig. 1E).

Besides these three types of interactions, a few encounters were observed where the opossum and the snake faced each other




FIG. 2. Scheme of attack and defense strategies in the encounters of opossums and snakes. Three different types of reaction are observed in the rattlesnakes after the first attack by the opossum. N = number of filmed events.

without moving for a long time. The two animals remained in this position until the snake slowly moved backwards.

During predation, the prey handling behavior was quite regular: the opossum remained in a sitting position while holding the snake with one of the forelimbs (Figs. 1F, 1G). One of the ends of the snake was introduced into the mouth laterally and was chewed with the lateral teeth (Figs. 1F, 1G). The other forelimb supported the animal (Figs. 1F, 1G). The forefeet were used alternately but for short periods of time were used together to tear apart the harder parts of the snake. Generally the whole snake was eaten uninterruptedly (Fig. 1H) with an average duration of 12 min.

The opossums killed and ate approximately 80% of the offered rattlesnakes. Taking into account the filmed captures where the attack was directed to the tail or to the head ($N = 22$), the opossums seemed to direct the attack significantly to the tail ($N = 16$; $p < 0.05$, $df = 1$). In three filmed captures the opossums directed the attack to the middle of the body.

During all experiments no observable effects of envenomation were recognized.

DISCUSSION

The way the opossums behaved while eating the snakes was very regular. The opossums were always in the same posture when observed. They used only one forelimb at a time to hold the snake and introduce it into the mouth. Ivanco et al. (1996) also verified the use of a single limb in another opossum species, *Monodelphis domestica*, when feeding or preying, and suggested that this behavior is fixed and species-typical.

During the time preceding the effective predation, when the encounter between opossum and rattlesnake took place, great behavioral variations were noted in both animals.

Before the attack by the opossum, the differences in reaction presented by the rat-

tlesnakes indicated that some times they were not able to notice the opossum, moving calmly around the tank or remaining quietly coiled. Most times, however, the rattlesnakes demonstrated by their behavior that they could recognize the opossum as a predator; in these cases some of the typical behaviors of the defensive escalation of the rattlesnakes were observed, such as immobility, coiling, cocking, and rattling. Among these defensive behaviors, the strategy of immobility in nature can be very valuable when associated with a cryptic coloration pattern and may constitute an efficient defense used by these animals against predators (Greene et al., 1978; Herzog and Drummond, 1984; Cloudsley-Thompson, 1994). Rattling many times occurred despite the snake not being able to see the opossum inside the box. Since it was not observed in any of the control experiments, rattling indicates that most times the snakes seemed to identify somehow the danger, possibly by chemical signs, as proposed by Weldon et al. (1992).

On a significant number of occasions the attack of the opossum was directed to the tail. That was observed more frequently when the rattlesnakes remained immobile, a strategy that at first view is difficult to interpret. On the other hand, during the approach, it was observed on a few occasions that the opossum, when noticing the snake coiling and preparing to strike, rapidly killed it by attacking and chewing its head. The same happened in the few cases when the rattlesnakes were able to bite the opossum even after being captured by the tail. In other cases, when the snake reacted to the attack by exhibiting erratic behavior, it was immobilized through bites along the whole body. The analysis of these different situations observed during the attack indicated that any type of active reaction presented by the snake caused an immediate fatal attack by the opossum. Although in our experimental conditions attacks directed to the tail did not prevent the snake from

being killed by the opossum, in nature they may confer an advantage to the snake, that by remaining immobile has a chance of escaping without being severely injured, as has been already observed for lizards by Greene et al. (1978). These data seem to be in accordance with Herzog and Burghardt (1974), who affirmed that for many predators, prey movement is a critical factor in mediating attack.

In contrast to our results indicating some preference of the opossums for capturing the snakes by the tail, Sazima (1992) reported that *D. marsupialis* when attacking *Bothrops jararaca* usually goes first to the head or neck region. In our experiments with *Crotalus durissus*, in many cases, *Didelphis* grasped the tail first, giving the snakes a chance to bite. This observation, at first view, seems to be contradictory since the predatory behavior of ophiophagous animals (mammals or birds) usually consists in attacking the head or the region just behind the head (Kaufmann and Kaufmann, 1965; Perez et al., 1978). However, ophiophagous animals such as *Conepatus sp.* and *Galictis sp.* have been observed attacking snakes at the tail (Ribeiro, 1940; Jackson, 1979).

It is possible that in our study the apparent preference of *D. marsupialis* for attacking the tail of *C. durissus* may be caused by an attraction of the opossums to cloacal odors of the rattlesnakes that can misdirect the attack. In nature, such attraction of the predator to the tail, which is a more disposable portion of the body, could help the prey to escape or counterattack (Greene, 1988; Alcock, 1993).

Two species of opossums of the genus *Didelphis* occur in Brazil: *D. albiventris* and *D. marsupialis* (Cerqueira, 1985). The former lives in open fields such as "cerrado" and "caatinga" and the latter is distributed in forests (Cerqueira, 1985; Emmons, 1990). On the other hand, *Crotalus durissus* is a species typical of open fields while *Bothrops jararaca* is distributed in forests

(Sazima, 1992; Campbell and Lamar, 1989). In this way, one would expect the opossums to have resistance only to snake venoms from the same habitat. In fact, Mous-satché et al. (1990) have demonstrated that *D. marsupialis* remains unharmed by *B. jararaca* venom. In addition, they mentioned that this marsupial has partial resistance to *Crotalus durissus* venom. On the basis of this information, our experiments aimed at comparing behavioral results with the biochemical data of Mous-satché et al. (1990). Although *D. marsupialis* and *C. durissus* are not sympatric in nature, we observed that, at least in captivity, predation was effective. The injection of venom by snakebites apparently did not affect the predation as has already been observed for other predators, including mammals and birds (Duvall et al., 1985).

In spite of the limitations imposed by a behavioral experiment conducted in captivity, the present observations strongly suggest that *D. marsupialis* is an effective snake predator in nature. This supposition is mainly based on the great interest and ability shown by *D. marsupialis* in capturing *Crotalus durissus*, which were comparable to the interest they showed when presented with different types of food. This idea is also reinforced by the great tolerance these animals showed to the snake venom.

ACKNOWLEDGEMENTS

The authors wish to thank Ms. Simone Jared, Mrs. Laurinda A. Soares, Mrs. Aláide M. Marques, Mr. Valdir J. Germano and Mr. Yutaka Kanasi, for their valuable technical assistance. We are also thankful to Dr. André Eterovic and Mr. Lucio Paraná Catani for suggestions. Fundação Butantan and FAPESP (proc. 98/8183-5) provided the financial support. Dr. Osvaldo Augusto Sant'Anna is supported, in part, by CNPq Brazil.

LITERATURE CITED

- ALCOCK, J. 1993. *Animal Behavior: An Evolutionary Approach*. Sinauer Associates, Sunderland, Massachusetts. 625 p.
- BRODIE, E. D. III. 1993. Differential avoidance of coral snake banded patterns by free-ranging avian predators in Costa Rica. *Evolution* 47: 227-235.
- CAMPBELL, J. A. AND W. W. LAMAR. 1989. *The Venomous Reptiles of Latin America*. Comstock Publishing Associates. Cornell Univ. Press, Ithaca, New York. 425 p.
- CERQUEIRA, R. 1985. The distribution of *Didelphis* in South America (Polyprotodontia, Didelphidae). *J. Biogeogr.* 12: 135-145.
- CLOUDSLEY-THOMPSON, J. L. 1994. *Predation and Defense among Reptiles*. R & A Publishing, Somerset, England. 138 p.
- CORDERO, G. A. R. AND R. A. B. NICOLAS. 1986. Feeding habits of the opossum (*Didelphis marsupialis*) in northern Venezuela. *Field. Zool.* 39: 125-131.
- DOMONT, G. B., J. PERALES, AND H. MOUSSATCHÉ. 1991. Natural anti-snake venom proteins. *Toxicon* 29: 1183-1194.
- DUVALL, D., M. B. KING, AND K. J. GUTZWILLER. 1985. Behavioral ecology and ethology of the prairie rattlesnake. *Natl. Geogr. Res.* 1: 80-111.
- EMMONS, L. H. 1990. *Neotropical Rainforest Mammals: A field Guide*. The University of Chicago, Chicago. 281 p.
- FITCH, H. S. 1960. Autoecology of the copperhead. *Univ. Kansas. Pub. Mus. Nat. Hist.* 13: 85-288.
- GREENE, H. W. 1973. Defensive tail display by snakes and amphisbaenians. *J. Herpetol.* 7: 143-161.
- GREENE, H. W. 1988. Antipredator mechanisms in reptiles. p. 1-152. *In*: C. Gans and R. B. Huey (eds.), *Biology of the Reptilia*, Vol. 16, Defense and Life History. Allan R. Liss, New York.
- GREENE, H. W., G. M. BURGHARDT, B. A. DUGAN, AND A. S. RAND. 1978. Predation and the defensive behavior of green iguanas (Reptilia, Lacertilia, Iguanidae). *J. Herpetol.* 12(2): 169-176.
- HERZOG, H. A. JR. AND G. M. BURGHARDT. 1974. Prey movement and predatory behavior of juvenile western yellow-bellied racers, *Coluber constrictor mormon*. *Herpetologica* 30: 285-289.
- HERZOG, H. A. JR. AND H. DRUMMOND. 1984. Tail autotomy inhibits tonic immobility in geckos. *Copeia* 1984 (3): 763-764.
- IVANCO, T. L., S. M. PELLIS, AND I. Q. WHISHAW. 1996. Skilled forelimb movements in prey catching and in reaching by rats (*Rattus norvegicus*) and opossums (*Monodelphis domestica*): relations to anatomical differences in motor systems. *Behav. Brain Res.* 79: 163-181.
- JACKSON, J. F. 1979. Effects of some ophidian tail displays on the predatory behavior of grison (*Galictis* sp.). *Copeia* 1979(1): 169-172.
- JARED, C., M. M. ANTONIAZZI, AND S. M. ALMEIDA-SANTOS. 1998. Predation of snakes by the young of opossum (*Didelphis marsupialis*) in captivity. *The Snake* 28: 68-70.
- KAUFMANN, J. H. AND A. KAUFMANN. 1965. Observations of the behavior of tayras and grisons. *Z. Säugetierkunde* 30: 146-155.
- MOUSSATCHÉ, H., A. YATES, F. LEONARDI, AND L. BORCHE. 1979. Mechanisms of resistance of the opossum to some snake venoms. *Toxicon* 17: 130.
- MOUSSATCHÉ, H., J. PERALES, A. G. C. N. FERREIRA, S. L. G. ROCHA, C. G. VILLELA, AND G. B. DOMONT. 1990. Resistência de marsupiais brasileiros aos venenos de *Bothrops jararaca* e *Crotalus durissus terrificus*. *Proc. Vth Ann. Meet. Fed. Soc. Exp. Biol., MG, Brazil*. p. 397.
- NEVES-FERREIRA, A. G., J. PERALES, M. OVALDIA, H. MOUSSATCHÉ, AND G. B. DOMONT. 1997. Inhibitory properties of the antithrotophic complex from the South American opossum (*Didelphis marsupialis*) serum. *Toxicon* 35: 849-863.
- PERALES, J., H. MOUSSATCHÉ, S. MARANGONI, B. OLIVEIRA, AND G. B. DOMONT. 1994. Isolation and partial characterization of an antithrotophic complex from serum of South American Didelphidae. *Toxicon* 32: 1237-1249.
- PEREZ, J. C., H. C. HAWS, V. E. GARCIA, AND B. M. JENNINGS, III. 1978. Resistance of warm-blooded animals to snake venoms. *Toxicon* 16: 375-383.
- PEREZ, J. C., S. PICHYANGKUL, AND V. E. GARCIA. 1979. The resistance of three species of warm-blooded animals to western diamond-back rattlesnake (*Crotalus atrox*) venom. *Toxicon* 17: 601-607.
- RIBEIRO, L. 1940. *Medicina no Brasil*. Imprensa Nacional, Rio de Janeiro. 409 p.

- SAZIMA, I. 1992. Natural history of the jararaca pitviper, *B. jararaca* in Southeastern Brazil. p. 199–216. In: J. A. Campbell and E. D. Brodie, Jr. (eds.), Biology of the Pitvipers. Selva Publ., Tyler, Texas.
- SILVA, M. JR. 1956. O Ofidismo no Brasil. Ministério da Saúde, Rio de Janeiro. 346 p.
- VELLARD, J. 1945. Resistencia de los "Didelphis" (Zarigueya) a los venenos ofidicos. Rev. Bras. Biol. 5: 463–467.
- WELDON, P. J., R. ORTIZ, AND T. R. SHARP. 1992. The Chemical Ecology of Crotalinae Snakes. p. 309–319. In: J. A. Campbell and E. D. Brodie, Jr. (eds.), Biology of the Pitvipers. Selva Publ., Tyler, Texas.
- WERNER, R. M. AND J. A. VICK. 1977. Resistance of the opossum (*Didelphis virginiana*) to envenomation by snakes of the family Crotalidae. Toxicon 15: 29–33.

Accepted: 23 December 1999